

Generally, air masses move the way they do because of the forces that act upon them.

• Definition: A force is something that pushes or pulls on an object, causing it to speed up, slow down, and/or change directions.




Generally, air masses move the way they do because of the forces that act upon them.

• Definition: A force is something that pushes or pulls on an object, causing it to speed up, slow down, and/or change directions.




Generally, air masses move the way they do because of the forces that act upon them.

• Definition: A force is something that pushes or pulls on an object, causing it to speed up, slow down, and/or change directions.



- Generally, air masses move the way they do because of the forces that act upon them.
- Definition: A force is something that pushes or pulls on an object, causing it to speed up, slow down, and/or change directions.


- To paraphrase Joe Biden, the development of Newton's laws was a big (censored) scientific deal!
  - Seen in the context of the time, Newton's laws of motion were perhaps the biggest single advance in the history of science
  - As never before, scientists could make quantitative,
     mathematical predictions for a wide range of physical systems
  - In meteorology, that meant predicting the weather. But of course that didn't happen overnight.....



1680's: Motivated by planetary orbits, Newton develops his fundamental laws of motion, along with the basic ideas of differential calculus. Simultaneously (and independently), Leibniz develops calculus in a form closer to that used today.




1750's: Leonard Euler develops a form of Newton's laws suitable for "perfect" (frictionless) fluids and gases.





1820's: Claude-Louis Navier and George Stokes extend Euler's equations to include the effects of friction. Essentially these same laws are used to predict the weather today.



1920's: Employing an army of people with calculators, Lewis Fry Richardson makes the first attempt (ultimately unsuccessful) to use Newton's laws to predict the weather.

Jon Von Neumann



1950's: The first modern electronic computer (the ENIAC) makes the first successful weather forecast using Newton's laws.


- So the key to understanding atmospheric motions is to understand the forces that produce these motions. And what, pray tell, are these forces you speak of?
- Well, for our purposes, the main ones are:

## The Pressure Gradient Force (PGF)

For any given mass of air, the air surrounding it exerts pressure forces across each of its sides.

 The net pressure force exerted by the surrounding air is called the pressure gradient force (or PGF)


 The PGF always pushes from higher pressure towards lower pressure



Higher pressure in Tank A causes water to flow towards Tank B


 On a surface map, the PGF is perpendicular to the isobars, pointing from higher pressure to lower pressure

 The strength of the PGF depends on the contour spacing:





More closely spaced contours imply a bigger PGF!


- On an upper-level chart,
   the PGF is perpendicular
   to height contours,
   pointing from higher to
   lower heights
- And as at the surface, the strength of the PGF depends on the contour spacing:





Again, tighter contours mean a stronger PGF!
























 To see how this works, consider an object moving in a straight line at constant speed, as seen from a coordinate system with fixed axes


> (Note that there's no force on the object, since it doesn't change speed or direction.)









































































 In our case, the moving object is any small mass of air as it moves through the atmosphere.
 And because the Earth is rotating, this air will appear to be deflected by a Coriolis force, much like the example shown.




# Two important properties of the Coriolis force:




An observer on the rotating Earth spins at different rates, depending on location:

 An observer at the North Pole spins counter-clockwise




An observer on the rotating Earth spins at different rates, depending on location:

- An observer at the North Pole spins counter-clockwise
- At the equator, the observer doesn't spin at all (he just moves forward)




An observer on the rotating Earth spins at different rates, depending on location:

- An observer at the North Pole spins counter-clockwise
- At the equator, the observer doesn't spin at all (he just moves forward)
- At the South Pole, the observer spins clockwise



An observer on the rotating Earth spins at different rates, depending on location:

- An observer at the North Pole spins counter-clockwise
- At the equator, the observer doesn't spin at all (he just moves forward)
- At the South Pole, the observer spins clockwise



From this, we can infer.....

- In between, the force varies with sin of the latitude  $\phi$ ; i.e.,

 The Coriolis force pulls in opposite directions for the Northern and Southern Hemispheres

Northern Hemisphere Southern Hemisphere