

Section 10.9: Taylor Polynomials



Definition: Let $f(x)$ be a function. Recall the Taylor Series for $f(x)$ at $x = a$ is

$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$. A partial sum of a Taylor Series is called a Taylor Polynomial. More specifically, the n^{th} degree **Taylor Polynomial** for $f(x)$ at $x = a$ is


$$T_n(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x - a)^i$$

where $f^{(i)}(a)$ is the i^{th} derivative of $f(x)$ at $x = a$.

EXAMPLE 1: Find the first and second degree Taylor Polynomial for $f(x) = \sqrt{x}$ at $x = 1$.




EXAMPLE 2: Find $T_1(x)$, $T_3(x)$, and $T_5(x)$ for $f(x) = \sin x$ at $x = 0$. Notice by viewing the graph below how the Taylor Polynomials better approximate $f(x)$ at $x = 0$ as n gets larger.



EXAMPLE 3: If $f(x) = x^{1/3}$, find $T_2(x)$ at $x = 8$.

Definition: The **Remainder** in using $T_n(x)$ to approximate $f(x)$ for x near a is defined to be $R_n(x) = f(x) - T_n(x)$. Thus it follows that $f(x) = T_n(x) + R_n(x)$.

Below, you will see a graph of the function and the Taylor Polynomial we found in example 3. Note that the function and the Taylor Polynomial are the same at $x = a$ (where $a = 8$ in this example), and as x deviates from $x = 8$, the Taylor Polynomial deviates from $f(x)$. Thus, the closer we are to a , the smaller the remainder.



Taylor's Inequality: An upper bound on the absolute value of the remainder is

$$|R_n(x)| \leq \frac{M}{(n+1)!} |x-a|^{n+1}$$

where $|f^{(n+1)}(x)| \leq M$ for x in an interval containing a .

EXAMPLE 4: If $f(x) = \ln x$ and $n = 3$, find $T_n(x)$ at $x = 4$. Then, use Taylor's inequality to find an upper bound on the remainder if $3 \leq x \leq 5$.

EXAMPLE 5: If $f(x) = \cos x$, $n = 2$. Find $T_n(x)$ at $x = \frac{\pi}{4}$. Then use Taylor's inequality to find an upper bound on the remainder if $\frac{\pi}{6} \leq x \leq \frac{2\pi}{3}$.

EXAMPLE 6: Recall $\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$ is the Maclaurin Series for $\sin x$.

(a) Using the Alternating Series Estimation Theorem, what is the maximum error possible in using the approximation

$$\sin x \approx x - \frac{x^3}{3!} + \frac{x^5}{5!} \text{ for } |x| < 0.3?$$

(b) Use $\sin x \approx x - \frac{x^3}{3!} + \frac{x^5}{5!}$ to approximate $\sin(1.2^\circ)$.

(c) Use the Alternating Series Estimation Theorem to estimate the range of values of x for which the given approximation is accurate to within the stated error.

$$\sin x \approx x - \frac{x^3}{3!} + \frac{x^5}{5!}, \text{ error} < 0.000001.$$