

Bounded Operators & Closed Subspaces

by

Francis J. Narcowich

November 2014¹

1 Bounded operators & examples

Let V and W be Banach spaces. We say that a linear transformation $L : V \rightarrow W$ is *bounded* if and only if there is a constant K such that $\|Lv\|_W \leq K\|v\|_V$ for all $v \in V$. Equivalently, L is bounded whenever

$$\|L\|_{op} := \sup_{v \neq 0} \frac{\|Lv\|_W}{\|v\|_V} \quad (1.1)$$

is finite. $\|L\|_{op}$ is called the norm of L . Frequently, the same operator may map another space $\tilde{V} \rightarrow \tilde{W}$, rather than $V \rightarrow W$. When this happens, we will need to note which spaces are involved. For instance, if V and W are the spaces involved, we will use the notation $\|L\|_{V \rightarrow W}$ for the operator norm. In addition to the expression given in (1.1), it is easy to show that $\|L\|_{op}$ is also given by

$$\|L\|_{op} := \min\{K > 0 : \|Lv\|_W \leq K\|v\|_V \ \forall v \in V\}. \quad (1.2)$$

As usual, we say $L : V \rightarrow W$ is continuous at $v \in V$ if and only if for every $\varepsilon > 0$ there is a $\delta > 0$ such that $\|Lu - Lv\|_W < \varepsilon$ whenever $\|u - v\|_V < \delta$. Of course, this is just the standard definition of continuity. Be aware that it holds whether or not L is linear. When L is linear, the distinction between u, v becomes irrelevant, because $\|Lu - Lv\|_W = \|L(u - v)\|_W$. From this it immediately follows that L will be continuous at every $v \in V$ whenever it is continuous at $v = 0$. The proposition below connects boundedness and continuity for linear transformations. The proof is left as an exercise.

Proposition 1. *A linear transformation $L : V \rightarrow W$ is continuous if and only if it is bounded.*

We will now provide a number of examples of bounded operators and unbounded operators.

¹Revised October 2019

Example 1. Let $L : C[0, 1] \rightarrow C[0, 1]$ be given by $Lu(x) = \int_0^1 k(x, y)u(y)dy$, where $k \in C(R)$, $R = [0, 1] \times [0, 1]$. We have that $|Lu(x)| \leq \int_0^1 |k(x, y)| |u(y)| dy$, so $|Lu(x)| \leq \|k\|_{C(R)} \|u\|_{C([0,1])}$. Consequently, $\|L\|_{C \rightarrow C} \leq \|k\|_{C(R)} \|u\|_{C([0,1])}$

Example 2. Hilbert-Schmidt operators.

Definition 1. Let $R = [0, 1] \times [0, 1]$ and let $k : R \rightarrow \mathbb{R}$. If $k \in L^2(R)$, then k is called a *Hilbert-Schmidt kernel*.

Proposition 2. Let k be a Hilbert-Schmidt kernel. The linear operator $Lu(x) = \int_0^1 k(x, y)u(y)dy$ maps $L^2[0, 1] \rightarrow L^2[0, 1]$ and is bounded. Moreover, $\|L\|_{L^2 \rightarrow L^2} \leq \|k\|_{L^2(R)}$.

Proof. Since $k(x, y) \in L^2(R)$, $\int_R |k(x, y)|^2 dx dy < \infty$, we have that $|k(x, y)|^2 \in L^1(R)$. Fubini's theorem then implies that $\int_0^1 |k(x, y)|^2 dy$ exists for almost every x and, in x , is in $L^1[0, 1]$. But this also implies that for almost every x , $|k(x, y)|^2$ is L^2 in y . Hence, by Schwarz's inequality,

$$|Lu(x)|^2 = \left| \int_0^1 k(x, y)u(y)dy \right|^2 \leq \int_0^1 |k(x, y)|^2 dy \underbrace{\int_0^1 |u(y)|^2 dy}_{\|u\|_{L^2}^2}.$$

Integrating both sides in x then yields $\|Lu\|_{L^2[0,1]}^2 \leq \|k\|_{L^2(R)}^2 \|u\|_{L^2[0,1]}^2$, so $\|Lu\|_{L^2[0,1]} \leq \|k\|_{L^2(R)} \|u\|_{L^2[0,1]}$. Then by (1.2), we see that $\|L\|_{L^2 \rightarrow L^2} \leq \|k\|_{L^2(R)}$, which completes the proof. \square

Example 3. Consider $L^2[0, 1]$. The differentiation operator $D = \frac{d}{dx}$ is defined on all $f \in C^1[0, 1]$, which is dense in L^2 because it contains the set of polynomials. The question is whether D is bounded, or at least can be extended to a bounded operator on L^2 . The answer is no. Let $u_n(x) := \sqrt{2} \sin(n\pi x)$. These functions are in $C^1[0, 1]$ and they satisfy $\|u_n\|_{L^2} = 1$. Since $Du_n = n\pi\sqrt{2} \cos(n\pi x)$, $\|Du_n\|_{L^2} = n\pi$. Consequently,

$$\frac{\|Du_n\|_{L^2}}{\|u_n\|_{L^2}} = n\pi \rightarrow \infty, \text{ as } n \rightarrow \infty.$$

Thus D is an *unbounded* operator on $L^2[0, 1]$.

The situation changes if we use a different space. Consider the Sobolev space $H^1[0, 1]$, which has the inner product

$$\langle f, g \rangle_{H^1} = \int_0^1 f(x)\overline{g(x)} + f'(x)\overline{g'(x)} dx.$$

The operator $D : H^1 \rightarrow L^2$ turns out to be bounded. In fact, one can show that $\|D\|_{H^1 \rightarrow L^2} = 1$. (It's easy to show that $\|D\|_{H^1 \rightarrow L^2}$ is at *most* 1. Showing that it's exactly one requires more work.)

2 Closed subspaces

The usual definition of subspace holds for Banach spaces and for Hilbert spaces. Such subspaces inherit norms and/or inner products from the larger spaces. They are said to be *closed* if they contain all of their limit points.

Finite dimensional subspaces are always closed. Earlier, when we discussed completeness of an orthonormal set $U = \{u_n\}_{n=1}^\infty$ in a Hilbert space \mathcal{H} , we saw that the space $\mathcal{H}_U = \{f \in \mathcal{H} : f = \sum_n \langle f, u_n \rangle u_n\}$ is closed in \mathcal{H} . When $C[0, 1]$ is considered to be a subspace of $L^2[0, 1]$, it is not closed. However, $C[0, 1]$ is a closed subspace of $L_\infty[0, 1]$.

Given a subspace V of a Hilbert space \mathcal{H} , we define the *orthogonal complement* of V to be

$$V^\perp := \{f \in \mathcal{H} : \langle f, g \rangle = 0 \ \forall g \in V\}.$$

Proposition 3. V^\perp is a closed subspace of \mathcal{H} .

Proof. Let $\{f_n\}_{n=1}^\infty$ be a sequence in V^\perp that converges to a function $f \in \mathcal{H}$. Since each f_n is in V^\perp , $\langle f_n, g \rangle = 0$ for every $g \in V$. Also, because the inner product is continuous, $\lim_{n \rightarrow \infty} \langle f_n, g \rangle = \langle f, g \rangle$. It immediately follows that $\langle f, g \rangle = 0$, so $f \in V^\perp$. Consequently, V^\perp is closed in \mathcal{H} . \square

Bounded linear operators mapping $V \rightarrow W$, where V and W are Banach spaces, have all of the usual subspaces associated with them. Let $L : V \rightarrow W$ be bounded and linear. The domain of L is $D(L) = V$. The range of L is defined as $R(L) := \{w \in W : \exists v \in V \text{ for which } Lv = w\}$. Finally, the null space (or kernel) of L is $N(L) := \{v \in V : Lv = 0\}$.

Proposition 4. If $L : V \rightarrow W$ is bounded and linear, then the null space $N(L)$ is a closed subspace of V .

Proof. The proof again relies on the continuity of L . If $\{f_n\}_{n=1}^\infty$ is a sequence in $N(L)$ that converges to $f \in V$. By Proposition 1, L is continuous, so $\lim_{n \rightarrow \infty} Lf_n = Lf$. But, because $f_n \in N(L)$, $Lf_n = 0$. Combining this with $\lim_{n \rightarrow \infty} Lf_n = Lf$, we see that $Lf = 0$ and so $f \in N(L)$. Thus, $N(L)$ is a closed subspace of V . \square

Previous: X-ray tomography and integral equations

Next: the projection theorem, Reisz representation theorem and the Fredholm alternative