

ON A DIRECTIONALLY REINFORCED RANDOM WALK

ARKA P. GHOSH, REZA RASTEGAR, AND ALEXANDER ROITERSHTEIN

ABSTRACT. We consider a generalized version of a directionally reinforced random walk, which was originally introduced by Mauldin, Monticino, and von Weizsäcker in [20]. Our main result is a stable limit theorem for the position of the random walk in higher dimensions. This extends a result of Horváth and Shao [13] that was previously obtained in dimension one only (however, in a more stringent functional form).

1. INTRODUCTION

In this paper we study the following *directionally reinforced random walk*. Fix $d \in \mathbb{N}$ and a finite set U of distinct unit vectors in \mathbb{R}^d (see Remark 2.8 at the end of Section 2 below, where a suitable alternative Markovian setup in a general state space is discussed). The vectors in U serve as feasible directions for the motion of the random walk. To avoid trivialities we assume that U contains at least two elements. Let $X_t \in \mathbb{R}^d$ denote the position of the random walk at time t . Throughout the paper we assume that $X_0 = 0$. The random walk changes its direction at random times

$$s_1 := 0 < s_2 < s_3 < s_4 < \dots$$

We assume that the time intervals

$$T_n := s_{n+1} - s_n, \quad n \in \mathbb{N},$$

are independent and identically distributed. Let $\eta_n \in U$ be the direction of the walk during time interval $[s_n, s_{n+1})$. We assume that $\eta := (\eta_n)_{n \geq 1}$ is an irreducible stationary Markov chain on U which is, furthermore, independent of $(s_n)_{n \in \mathbb{N}}$.

For $t > 0$, let $N_t := \sup\{k \geq 1 : s_k \leq t\}$ be the number of times that the walker changes direction before time $t > 0$. Then

$$(1.1) \quad X_t = \sum_{i=1}^{N_t-1} \eta_i T_i + (t - s_{N_t}) \eta_{N_t}.$$

Notice that $N_t \geq 1$ with probability one, due to the convention $s_1 = 0$ that we have made.

The continuous time random walk $(X_t)_{t \geq 0}$ defined above is essentially the model introduced by Mauldin, Monticino, and von Weizsäcker in [20] and further studied by Horváth and Shao in [13] and by Siegmund-Schultze and von Weizsäcker in [25]. The technical difference between our model and the variant which has been studied in [13] is that in the latter, the next direction of the motion is chosen uniformly

Received by the editors May 31, 2012; Revised September 18, 2012.

2010 *Mathematics Subject Classification*. Primary: 60F05, 60F15; Secondary: 60J25, 70B05.

©XXXX American Mathematical Society

from the available set of “fresh directions”, while we do not impose any restrictions on the transition kernel of η besides irreducibility.

The original model proposed in [20] was inspired by certain phenomena that occur in ocean surface waves (cf. [30]) and was designed to reproduce the same features within a probabilistic framework. The main topic of [20] and [25] is recurrence-transience criteria. Horváth and Shao in [13] studied scaling limits of the random walk in different regimes, answering some of the questions which have been posed in [20].

We remark that somewhat related random walk models have been considered by Allaart and Monticino in [1, 2] and by Gruber and Schweizer in [12]. In the context of random walks in random environment, a similar in spirit model of *persistent* random walks was introduced by Szász and Tóth in [28, 29]. The common feature of “generic versions” of all models mentioned above is that the underlying random motion has a tendency to persist in its current direction.

Closely related to persistent random walks are recurrent “random flights” models where changes of the direction of the random motion follow a Poisson random clock. These models can be traced back to Pearson’s random walk [10, 15] and Goldstein-Kac one-dimensional “telegraph process” [16, 23]. Random flights have been intensively studied since the introduction of the telegraph process in the early 50’s, see for instance [11, 17, 18, 19, 22, 26] and references therein for a representative sample. An introductory part of [18] provides a short authoritative and up to a date survey of the field. We remark that, somewhat in contrary to directionally reinforced random walks, the main focus of the research in this area is on finding explicit form of limiting distributions for these processes.

In statistical physics, the model (1.1) with i.i.d. jump vectors η_i is a certain kind of *coupled continuous time random walk*. Limit theorems for coupled continuous time random walks have been developed in [5, 14, 21, 27], focusing on the case where the waiting times have infinite mean.

The main goal of this paper is to prove stable limit theorems for the directionally reinforced random walk in arbitrary dimension $d \geq 1$. In addition, we extend some limit results of [13] to our setting and also complement them by suitable laws of iterated logarithm. Our proofs can be easily carried over to a setup where the set of feasible directions U is not finite, but is rather supported (under the stationary law of the process) on a general Borel subset of the unit sphere; see Remark 2.8 below for more details.

Our results are stated in Section 2 whereas the proofs are contained in Section 3. The non-Gaussian limit theorems for the position of the random walk in higher dimensions, stated in Theorems 2.5 and 2.6, constitute the main contribution of this paper.

2. STATEMENT OF MAIN RESULTS

We first introduce a few notations. For a vector $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ let $\|x\| = \max_i |x_i|$. For (possibly, random) functions $f, g : \mathbb{R}_+$ (or \mathbb{N}) $\rightarrow \mathbb{R}$, write $f \sim g$ and $f(t) = o(g(t))$ to indicate that, respectively, $\lim_{t \rightarrow \infty} f(t)/g(t) = 1$ and $\lim_{t \rightarrow \infty} f(t)/g(t) = 0$, a. s. Let $\pi = (\pi_v)_{v \in U} \in \mathbb{R}^{|U|}$ be the unique stationary distribution of the Markov chain η and let

$$(2.1) \quad \mu = \sum_{v \in U} \pi_v v.$$

Thus $\mu = E(\eta_n) \in \mathbb{R}^d$ for each $n \in \mathbb{N}$.

The following theorem shows that a strong law of large numbers holds for X_t and that, under suitable second moment condition, the sample paths of the random walk are uniformly close to the sample paths of a drifted Brownian motion. We have:

Theorem 2.1.

(a) Suppose that $E(T_1^p) < \infty$ for some constant $p \in (1, 2)$. Then,

$$\|X_t - \mu t\| = o(t^{1/p}).$$

(b) If $E(T_1^p) < \infty$ for some constant $p > 2$, then (in an enlarged, if needed, probability space) there exist a process $\widehat{X} = (\widehat{X}_t)_{t \geq 0}$ distributed as X and a Brownian motion $(W_t)_{t \geq 0}$ in \mathbb{R}^d , such that,

$$\sup_{0 \leq t \leq T} \|\widehat{X}_t - \mu t - W_t\| = o(T^{1/p}).$$

Remark 2.2. The results stated in Theorem 2.1 as well as in Theorem 2.3 below are essentially due to [13]. In fact, the original proofs can be adapted to our more general setup. However, the proofs we give in Section 3 are shorter and somewhat simpler than the original ones. Furthermore, our proves can easily be seen working for the general Markov chain setup described in Remark 2.8 below.

The second part of Theorem 2.1 implies the invariance principle for $(X_{nt} - \mu nt)$ with the usual normalization \sqrt{n} . We next state an invariance principle and the corresponding law of iterated logarithm under a slightly more relaxed moment condition. Let $D(\mathbb{R}^d)$ denote the set of \mathbb{R}^d -valued càdlàg functions on $[0, 1]$ equipped with the Skorokhod J_1 -topology. We use notation \Rightarrow to denote the weak convergence in $D(\mathbb{R}^d)$. We have:

Theorem 2.3. For $n \in \mathbb{N}$, define a process S_n in $D(\mathbb{R}^d)$ by setting

$$(2.2) \quad S_n(t) = \frac{X_{nt} - \mu nt}{\sqrt{n}}, \quad t \in [0, 1].$$

If $E(T_1^2) < \infty$, then

(a) $S_n \Rightarrow W$, where $W = (W_t)_{t \geq 0}$ is a (possibly degenerate, but not identically equal to zero) d -dimensional Brownian motion.

(b) For every $x \in \text{Span}(U) \subset \mathbb{R}^d$, there is a constant $K(x) \in (0, \infty)$ such that

$$\limsup_{t \rightarrow \infty} \frac{(X_t - \mu t) \cdot x}{\sqrt{t \ln \ln t}} = K(x).$$

Furthermore, a similar statement holds for the \liminf .

We next consider the case when $E(T_1^2) = \infty$ and T_1 is in the domain of attraction of a stable law. Namely, for the rest of our results we impose the following assumption. Recall that a function $h : \mathbb{R}_+ \rightarrow \mathbb{R}$ is said to be *regularly varying of index* $\alpha \in \mathbb{R}$ if $h(t) = t^\alpha L(t)$ for some $L : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that $L(\lambda t) \sim L(t)$ for all $\lambda > 0$. We will denote the set of all regularly varying functions of index α by \mathcal{R}_α .

Assumption 2.4. For some $\alpha \in (0, 2]$ there exists a function $h \in \mathcal{R}_\alpha$ such that $\lim_{t \rightarrow \infty} h(t) \cdot P(T_1 > t) \in (0, \infty)$.

For $t > 0$ let

$$(2.3) \quad a_t = \begin{cases} \inf \{s > 0 : t \cdot P(T_1 > s) \leq 1\} & \text{if } \alpha < 2, \\ \inf \{s > 0 : ts^{-2} \cdot E(T_1^2; T_1 \leq s) \leq 1\} & \text{if } \alpha = 2 \end{cases}$$

If $h(t) \in \mathcal{R}_\alpha$ with $\alpha \in (1, 2]$ (and hence $E(T_1) < \infty$), one can obtain the following analogue of Theorem 2.3. It turns out that also in this case the functional limit theorem and the law of iterated logarithm for X_t inherit the structure of the corresponding statements for the partial sums of i.i.d. variables $\sum_{k=1}^n T_k$.

Theorem 2.5. *Let Assumption 2.4 hold with $\alpha \in (1, 2]$. Let*

$$S_t := \frac{X_t - \mu t}{a_t}, \quad t > 0.$$

We have:

(a) *If $\alpha \in (1, 2)$, then*

- (i) *S_t converges weakly to a non-degenerate multivariate stable law in \mathbb{R}^d .*
- (ii) *For every $x \in \text{Span}(U) \subset \mathbb{R}^d$ such that $x \cdot u > 0$ for some $u \in U$,*

$$\limsup_{t \rightarrow \infty} \frac{(X_t - \mu t) \cdot x}{a_t \cdot (\ln t)^{1/\alpha + \varepsilon}} = \begin{cases} 0 & \text{if } \varepsilon > 0, \\ \infty & \text{if } \varepsilon < 0 \end{cases} \quad \text{a. s.}$$

In particular,

$$\limsup_{t \rightarrow \infty} \left\{ \frac{(X_t - \mu t) \cdot x}{a_t} \right\}^{1/\ln \ln t} = e^{1/\alpha} \quad \text{a. s.}$$

(b) *If $\alpha = 2$ and $E(T_1^2) = \infty$, then S_t converges weakly to a non-degenerate multivariate Gaussian distribution in \mathbb{R}^d .*

For $\alpha \in (0, 1)$ we have the following limit theorem.

Theorem 2.6. *Let Assumption 2.4 hold with $\alpha \in (0, 1)$. Then $\frac{X_t}{t}$ converges weakly in \mathbb{R}^d to a non-degenerate limit.*

Remark 2.7. The limiting random law in the statement of Theorem 2.6 is specified in (3.24) below. The stable limit laws for X_t stated in Theorems 2.5 and Theorem 2.6 are extensions of corresponding one-dimensional results in [13]. The latter however are obtained in [13] in a more stringent functional form. The law of iterated logarithm given in Theorem 2.5 appears to be new even for $d = 1$.

Remark 2.8. Recall Markov chain $\eta = (\eta_n)_{n \geq 0}$ which records successive directions of the random walk. Let \mathcal{S}^{d-1} denote the d -dimensional unit sphere and let \mathcal{T}_d denote the σ -algebra of the Borel sets of \mathcal{S}^{d-1} . Denote by $H(x, A)$ transition kernel of η on $(\mathcal{S}^{d-1}, \mathcal{T}_d)$. We remark that

- (i) All the results stated in this section remain true for an arbitrary (not stationary) initial distribution of the Markov chain η .
- (ii) The proofs of our results given in Section 3 rest on the exploiting of a regenerative (renewal) structure associated with η , i.e. on the use of random times τ_n which are introduced below in Section 3.1. It is then not hard to verify that all the results stated in this section, with the only exception of the generalized law of iterated logarithm given in part (a)-(ii) of Theorem 2.5, remain true for a class of regenerative (in the sense of [3]) Markov chains η whose stationary distribution are supported on general Borel subsets of \mathcal{S}^{d-1} rather than on a finite set $U \subset \mathcal{S}^{d-1}$.

For instance, the following strong version of the classical Doeblin's conditions is sufficient for our purposes:

- There exist a constant $c_r > 1$ and a probability measure ψ on $(\mathcal{S}^{d-1}, \mathcal{T}_d)$ such that

$$(2.4) \quad c_r^{-1}\psi(A) < H(x, A) < c_r\psi(A) \quad \forall x \in \mathcal{S}, A \in \mathcal{T}_d.$$

A regenerative (renewal) structure for Markov chains which satisfies Doeblin's condition is described in [3]. Due to the fact that under the assumption (2.4), the kernel $H(x, A)$ is dominated uniformly from above and below by a probability measure ψ , such Markov chains share two key features with finite-state Markov chains. Namely, 1) the exponential bound stated in (3.2) holds for the renewal times which are defined in [3]; and 2) $c_r^{-1} < P_x(A)/P_y(A) < c_r$ for any non-null event $A \in \mathcal{T}_d$ and almost every states $x, y \in \mathcal{S}^{d-1}$ (with respect to the stationary law). Here P_x stands for the law of the Markov chain η starting from the initial state $x \in \mathcal{S}^{d-1}$. Once these two crucial properties are verified, our proofs (except only the proof of part (a)-(ii) of Theorem 2.5) work nearly verbatim for directionally reinforced random walks governed by a Markov chain η which satisfies condition (2.4).

3. PROOFS

This section is devoted to the proof of the results stated in Section 2 above. Some preliminary observations are stated in Section 3.1 below. The proof of Theorem 2.1 is contained in Section 3.2. Theorems 2.3 and 2.5 are proved in Section 3.3 and Section 3.4, respectively. Finally, the proof of Theorem 2.6 is given in Section 3.5.

3.1. Preliminaries. Our approach relies on the use of a renewal structure which is induced on the paths of the random walk by the cycles of the underlying Markov chain η . To define the renewal structure, set $\tau_0 = 0$ and let

$$\tau_{i+1} = \inf\{j > \tau_i : \eta_j = u_1\}, \quad i \geq 0.$$

Thus, for $i \geq 1$, τ_i are steps when the Markov chain η visits the distinguished state u_1 . Correspondingly, s_{τ_i} are successive times when the random walk chooses u_1 as the direction of its motion. Recall N_t from Section 1 (see a few lines preceding (1.1)). Denote by $c(t)$ the number of times that the walker chooses direction u_1 before time $t > 0$. That is,

$$c(t) := \sup\{i \geq 0 : s_{\tau_i} \leq t\} = \sum_{j=1}^{N_t} \mathbf{1}_{\{\eta_j = u_1\}},$$

where $\mathbf{1}_A$ stands for the indicator function of an event A . Notice that N_t is the unique mapping from \mathbb{R}_+ to \mathbb{Z}_+ which has the following property:

$$s_{N_t} \leq t < s_{N_t+1} \quad \text{and} \quad \tau_{c(t)} \leq N_t < \tau_{c(t)+1}.$$

For $i \geq 0$, let $\xi_i = \sum_{j=\tau_i+1}^{\tau_{i+1}} T_j \eta_j$. Then

$$(3.1) \quad X_t = \xi_0 + \sum_{i=1}^{c(t)-1} \xi_i + \sum_{j=\tau_{c(t)}+1}^{N_t} T_j \eta_j + (t - s_{N_t}) \cdot \eta_{N_t}.$$

The strong Markov property implies that the pairs $(\xi_i, \tau_{i+1} - \tau_i)_{i \in \mathbb{N}}$ form an i.i.d. sequence which is independent of (ξ_0, τ_1) . Furthermore, since η is an irreducible

finite-state Markov chain, there exist positive constants $K_1, K_2 > 0$ such that the inequality

$$(3.2) \quad P(\tau_{i+1} - \tau_i > t) \leq K_1 e^{-K_2 t}$$

holds uniformly for all reals $t \geq 0$ and all integers $i \geq 0$.

We next list some direct consequences of the law of large numbers that will be frequently exploited in the subsequent proofs. Let $v(n)$ be the number of times that the Markov chain η visits u_1 during its first n steps. Thus, while $c(t)$ is the number of visits of η to u_1 up to time $t > 0$ on the clock of the random walk, $v(n)$ is the number of occurrences of u_1 among first n directions of the random walk. In particular, $v(N_t) = c(t)$. Taking into account (3.2), the law of large numbers and the renewal theorem imply that

$$\lim_{n \rightarrow \infty} \frac{\tau_n}{n} = \lim_{n \rightarrow \infty} \frac{n}{v(n)} = E(\tau_2 - \tau_1) = \pi_1^{-1}, \quad \text{a. s.},$$

and, letting $\Lambda_k := \sum_{i=\tau_k+1}^{\tau_{k+1}} \eta_i$,

$$\mu = \lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n \eta_i}{n} = \lim_{n \rightarrow \infty} \frac{\sum_{k=0}^{v(n)} \Lambda_k}{n} = \pi_1 \cdot E(\Lambda_1), \quad \text{a. s.}$$

Since η and $(T_k)_{k \in \mathbb{N}}$ are independent, it follows that

$$(3.3) \quad E(\xi_1) = E(T_1) \cdot E(\Lambda_1) = \pi_1^{-1} \mu \cdot E(T_1).$$

Finally, $\frac{c(t)}{t} = \frac{v(N_t)}{t} = \frac{v(N_t)}{N_t} \cdot \frac{N_t}{t}$ yields

$$(3.4) \quad \lim_{t \rightarrow \infty} \frac{c(t)}{t} = \frac{\pi_1}{E(T_1)}, \quad \text{a. s.}$$

We now turn to the proofs of our main results.

3.2. Proof of Theorem 2.1.

Part (a) of Theorem 2.1. Recall (3.2) and observe that the moment condition of the theorem along with the independence of the Markov chain η and $(T_k)_{k \in \mathbb{N}}$ of each other, implies that

$$(3.5) \quad \begin{aligned} E(\|\xi_1\|^p) &\leq E[(s_{\tau_2} - s_{\tau_1})^p] = \sum_{n=1}^{\infty} P(\tau_2 - \tau_1 = n) \cdot E\left[\left(\sum_{k=1}^n T_k\right)^p\right] \\ &\leq K_1 \sum_{n=1}^{\infty} e^{-K_2(n-1)} n^p E(T_1^p) < \infty, \end{aligned}$$

where we used Minkowski's inequality and (3.2). It follows that $\|\xi_k\| = o(k^{1/p})$. Indeed, for any $\varepsilon > 0$, Chebyshev's inequality implies that

$$\sum_{k=1}^{\infty} P(\|\xi_k\| > k^{\frac{1}{p}} \varepsilon) = \sum_{k=1}^{\infty} P(\|\xi_k\|^p > \varepsilon^p k) \leq \varepsilon^{-p} E(\|\xi_1\|^p) < \infty,$$

and hence $P(\|\xi_k\| > k^{\frac{1}{p}} \varepsilon \text{ i. o.}) = 0$ by the Borel-Cantelli lemma.

For now we will make a simplifying assumption (to be removed later on) that $\mu = 0$. By virtue of (3.4), the Marcinkiewicz-Zigmund law of large numbers implies that

$$\lim_{t \rightarrow \infty} \frac{\sum_{i=0}^{c(t)-1} \xi_i}{t^{1/p}} = 0, \quad \text{a. s.}$$

Furthermore, by (3.1), $\|X_t - \sum_{i=0}^{c(t)-1} \xi_i\| \leq r_{c(t)}$, where

$$(3.6) \quad r_k := \sum_{i=\tau_k+1}^{\tau_{k+1}} T_i.$$

An argument similar to the one which we used to estimate the order of $\|\xi_n\|$, shows that with probability one $r_n = o(n^{1/p})$. Then (3.4) implies that

$$(3.7) \quad r_{c(t)} = o(t^{1/p}).$$

This completes the proof of part (a) of Theorem 2.1 for the particular case $\mu = 0$.

We now turn to the general case of arbitrary finite $\mu \in \mathbb{R}^d$. Let

$$(3.8) \quad \tilde{\eta}_i = \eta_i - \mu \quad \text{and} \quad \tilde{X}_t = \sum_{i=0}^{N_t} T_i \tilde{\eta}_i + (t - s_{N_t}) \tilde{\eta}_{N_t}.$$

Then \tilde{X}_t is a directionally reinforced random walk associated with $(T_n)_{n \in \mathbb{N}}$ and $\tilde{\eta} = (\tilde{\eta}_i)_{i \in \mathbb{N}}$. Since $E(\tilde{\eta}_i) = 0$, we have $\|\tilde{X}_t\| = o(t^{1/p})$. To complete the proof of part (a) of the theorem, observe that $X_t - \tilde{X}_t = \mu \cdot \sum_{i=1}^{N_t} T_i + \mu \cdot (t - s_{N_t}) = \mu t$. \square

Part (b) of Theorem 2.1. Recall (3.3). Let

$$\bar{\xi}_k := \xi_k - E(\xi_1) = \xi_k - E(T_1)\pi_1^{-1}\mu$$

and

$$\Delta_k := s_{\tau_{k+1}-1} - s_{\tau_k-1} - E(T_1)\pi_1^{-1}.$$

Let $\gamma_k = (\bar{\xi}_k, \Delta_k) \in \mathbb{R}^{d+1}$. Then $(\gamma_k)_{k \geq 1}$ is an i.i.d. sequence with $E(\gamma_1) = 0 \in \mathbb{R}^{d+1}$. Define

$$\Gamma(t) = \sum_{1 \leq k \leq t} \gamma_k.$$

By virtue of Theorem 1.2.1 in [9], there is a Brownian motion $(B(t))_{t \geq 0}$ in \mathbb{R}^{d+1} such that

$$\sup_{0 \leq t \leq T} \|\Gamma(t) - B(t)\| = o(T^{1/p}).$$

Then Theorem 2.3.6 in [9] implies that there exists a Brownian motion $(W(t))_{t \geq 0}$, such that

$$\sup_{0 \leq t \leq T} \left\| \sum_{k=0}^{c(t)-1} \xi_k - t\mu - W(t) \right\| = o(T^{1/p}).$$

Recall r_k from (3.6). Since

$$\sup_{0 \leq t \leq T} \left\| X_t - \sum_{k=0}^{c(t)-1} \xi_k \right\| \leq \sup_{0 \leq t \leq T} r_{c(t)},$$

it suffices to show that

$$(3.9) \quad \sup_{0 \leq t \leq T} r_{c(t)} = o(T^{1/p}).$$

Notice that

$$\sup_{0 \leq t \leq T} r_{c(t)} = \sup_{0 \leq k \leq c(T)} r_k.$$

Therefore, by virtue of (3.4), it suffices to show that

$$(3.10) \quad \lim_{n \rightarrow \infty} n^{-1/p} \cdot \sup_{0 \leq k \leq n} r_k = 0, \quad \text{a. s.}$$

Toward this end, let

$$g(n) = \max\{k \leq n : r_k \geq r_i \text{ for all } 1 \leq i \leq n\}, \quad n \in \mathbb{N}.$$

Thus $g(n) \leq n$ and $\sup_{0 \leq k \leq n} r_k = r_{g(n)}$. Furthermore, since r_k are i.i.d. random variables, $\lim_{n \rightarrow \infty} g(n) = \infty$ with probability one. Therefore, $r_n = o(n^{1/p})$ yields (3.10). The proof of Theorem 2.1 is completed. \square

3.3. Proof of Theorem 2.3.

Part (a) of Theorem 2.3. By (3.5), $E(\|\xi_1\|^2) < \infty$ under the conditions of the theorem. Assume first that $\mu = 0$. Then the invariance principle for i.i.d. sequences implies that

$$\frac{\sum_{k=1}^{\lfloor nt \rfloor} \xi_k}{\sqrt{n}} \Rightarrow W(t), \quad t \in [0, 1],$$

where $W(t)$ is a d -dimensional Brownian motion. It follows then from (3.4) and Theorem 14.4 in [6, p. 152] that

$$(3.11) \quad \frac{\sum_{k=0}^{c(nt)-1} \xi_k}{\sqrt{n}} \Rightarrow \sqrt{b} \cdot W(t), \quad t \in [0, 1],$$

where $b = \frac{\pi_1}{E(T_1)}$. Under the moment condition of Theorem 2.3 we have the following counterpart of (3.9):

$$\sup_{0 \leq t \leq T} r_{c(t)} = o(T^{1/2}).$$

Since $\|X_{nt} - \sum_{k=0}^{c(nt)-1} \xi_k\|$ is bounded above by $r_{c(nt)}$, it follows that

$$n^{-1/2} \cdot \left\| X_{nt} - \sum_{k=0}^{c(nt)-1} \xi_k \right\| \Rightarrow 0,$$

which implies the desired convergence of $n^{-1/2} \cdot X_{nt}$ when $\mu = 0$. To prove the general case of arbitrary $\mu \in \mathbb{R}^d$ one can apply the result with $\mu = 0$ to the Markov chain $\tilde{\eta}_n$ and the random walk \tilde{X}_t that were introduced in (3.8). The proof of part (a) of the theorem is completed. \square

Part (b) of Theorem 2.3. Suppose first that $\mu = 0$. For $x \in \text{Span}(U) \subset \mathbb{R}^d$ and $i \in \mathbb{N}$ define

$$(3.12) \quad \xi_{i,x} := \xi_i \cdot x.$$

Then, in view of (3.4), the law of iterated logarithm for i.i.d. sequences implies that there exists a constant $K(x) \in (0, \infty)$ such that

$$\limsup_{t \rightarrow \infty} \frac{\sum_{i=0}^{c(t)-1} \xi_{i,x}}{\sqrt{t \ln \ln t}} = K(x), \quad \text{a. s.}$$

By (3.3) and (3.7)

$$\lim_{t \rightarrow \infty} \frac{|X_t \cdot x - \sum_{i=0}^{c(t)-1} \xi_{i,x}|}{\sqrt{t \ln \ln t}} = 0, \quad \text{a. s.}$$

Thus

$$\limsup_{t \rightarrow \infty} \frac{X_t \cdot x}{\sqrt{t \ln \ln t}} = K(x), \quad \text{a. s.,}$$

in the case $\mu = 0$. To obtain the general case with an arbitrary $\mu \in \mathbb{R}^d$, apply this result to the random walk \tilde{X}_t defined in (3.8) and recall that $X_t - \tilde{X}_t = \mu t$. The proof of part (b) of Theorem 2.3 is completed. \square

3.4. Proof of Theorem 2.5.

Part (a)-(i) and part (b) of Theorem 2.5. Let $\overline{\mathbb{R}}_0^d := [-\infty, \infty]^d \setminus \{0\}$, where 0 stands for the zero vector in \mathbb{R}^d , and equip $\overline{\mathbb{R}}_0^d$ with the topology inherited from \mathbb{R}^d . Recall (see for instance [4, 24]) that a random vector $\xi \in \mathbb{R}^d$ is said to be regularly varying with index $\alpha > 0$ if there exists a function $a : \mathbb{R}_+ \rightarrow \mathbb{R}$, regularly varying with index $1/\alpha$, and a Radon measure ν_ξ on $\overline{\mathbb{R}}_0^d$ such that

$$(3.13) \quad nP(a_n^{-1}\xi \in \cdot) \xrightarrow{v} \nu_\xi(\cdot), \quad \text{as } n \rightarrow \infty,$$

where $a_n := a(n)$ and \xrightarrow{v} denotes the vague convergence of measures. We will denote by $\mathcal{R}_{d,\alpha,a}$ the set of all random d -vectors regularly varying with index α , associated with a given function $a \in \mathcal{R}_{1/\alpha}$ by (3.13). The measure ν is referred to as the *measure of regular variation* associated with ξ . We will also use the following equivalent definition of the regular variation for random vectors (see, for instance, [4, 24]). Let S^{d-1} denote the unit sphere in \mathbb{R}^d with respect to the norm $\|\cdot\|$. Then $\xi \in \mathcal{R}_{d,\alpha,a}$ if and only if there exists a finite Borel measure \mathfrak{S}_ξ on S^{d-1} such that for all $t > 0$,

$$(3.14) \quad nP(\|\xi\| > ta_n; \xi/\|\xi\| \in \cdot) \xrightarrow{v} t^{-\alpha} \mathfrak{S}_\xi(\cdot), \quad \text{as } n \rightarrow \infty,$$

where \xrightarrow{v} denotes the vague convergence of measures on S^{d-1} . The following well-known result is the key to the proof of the next lemma: If $\xi, \eta \in \mathcal{R}_{d,\alpha,a}$ and ξ, η are independent of each other, then $\xi + \eta \in \mathcal{R}_{d,\alpha,a}$. Furthermore, $\nu_{\xi+\eta} = \nu_\xi + \nu_\eta$ and $\mathfrak{S}_{\xi+\eta} = \mathfrak{S}_\xi + \mathfrak{S}_\eta$ (see, for instance, Equation 1.15 and Lemma 2.6 in [8]).

We have:

Lemma 3.1. *Let Assumption 2.4 hold. For $t \geq 0$, let a_t be defined as in (2.3).*

Then

- (a) $\sum_{\tau_1+1}^{\tau_2} T_i \in \mathcal{R}_{1,\alpha,a}$.
- (b) $\xi_1 \in \mathcal{R}_{d,\alpha,a}$.

Proof of Lemma 3.1. The claim of part (a) can be formally deduced from that of part (b). Thus we will focus on proving the more general claim (b).

First, observe that (3.14) implies that $T_1 u \in \mathcal{R}_{d,\alpha,a}$ for any $u \in U$. Let

$$H(u, v) = P(\eta_{n+1} = v | \eta_n = u), \quad u, v \in U,$$

be the transition matrix of the Markov chain η . Further, define a sub-Markovian kernel Θ by setting

$$\Theta(u, v) = H(u, v) \cdot \mathbf{1}_{\{v \neq u_1\}}, \quad u, v \in U.$$

Fix any $t > 0$ and a Borel set $B \subset S^{d-1}$, and let

$$A_n = \{\|\xi_1\| > ta_n; \xi_1/\|\xi_1\| \in B\}, \quad n \in \mathbb{N}.$$

Then,

$$\begin{aligned} P(\xi_1 \in A_n) &= \sum_{k=1}^{\infty} P(\tau_2 - \tau_1 = k) \\ &\quad \times P(T_1 u_1 + T_2 \eta_2 + \dots + T_k \eta_k \in A_n | \tau_2 - \tau_1 = k) \\ &= \sum_{k=1}^{\infty} \sum_{v_2 \neq u_1} \dots \sum_{v_k \neq u_1} \Theta(u_1, v_1) \dots \Theta(v_{k-1}, v_k) H(v_k, u_1) \\ &\quad \times P(T_1 u_1 + T_2 v_2 + \dots + T_k v_k \in A_n), \end{aligned}$$

where we assume that the sums $\sum_{v_2 \neq u_1} \dots \sum_{v_k \neq u_1}$ are empty if $k = 1$. Let

$$J_n(v_2, \dots, v_k) = T_1 u_1 + T_2 v_2 + \dots + T_k v_k.$$

Notice that for any $k \in \mathbb{N}$ and fixed set of vectors $v_2, \dots, v_k \in U$, we have

$$\begin{aligned} n \cdot P(J_n(v_2, \dots, v_k) \in A_n) &\leq n \cdot P(\|J_n(v_2, \dots, v_k)\| \geq ta_n) \\ &\leq nP\left(\sum_{j=1}^k T_j \geq ta_n\right) \leq nkP(T_1 \geq ta_n/k) \leq Ct^{-\alpha} k^{1+\alpha} \end{aligned}$$

for some $C > 0$. Furthermore,

$$\lim_{n \rightarrow \infty} n \cdot P(J_n(v_2, \dots, v_k) \in A_n) = t^{-\alpha} \left(\mathfrak{S}_{T_1 u_1}(B) + \sum_{j=2}^k \mathfrak{S}_{T_1 v_j}(B) \right).$$

Observe that the spectral radius of the matrix Θ is strictly less than one and that $\mathfrak{S}_{T_1 v_j}(B)$ is uniformly bounded from above by $\max_{v \in U} \mathfrak{S}_{T_1 v}(S^{d-1})$. Therefore, the dominated convergence theorem implies that the following limit exists and the identity holds:

$$\begin{aligned} &\lim_{n \rightarrow \infty} n \cdot P(\xi_1 \in A_n) \\ &= \sum_{k=1}^{\infty} t^{-\alpha} \sum_{v_2 \neq u_1} \dots \sum_{v_k \neq u_1} \Theta(u_1, v_1) \dots \Theta(v_{k-1}, v_k) H(v_k, u_1) \\ &\quad \times \left(\mathfrak{S}_{T_1 u_1}(B) + \sum_{j=2}^k \mathfrak{S}_{T_1 v_j}(B) \right). \end{aligned}$$

Since the spectral radius of Θ is strictly less than one, Fubini's theorem implies that the right-hand side of the above identity defines a measure on S^{d-1} . The proof of the lemma is therefore completed. \square

We are now in a position to complete the proof of the limit results stated in parts (a) and (b) of Theorem 2.5. Suppose first that $\mu = 0$. It follows from Lemma 3.1 and the stable limit theorem for i.i.d. sequences (see, for instance, Section 1.6 in [7, p. 75]) that

$$(3.15) \quad \frac{\sum_{k=1}^{\lfloor nt \rfloor} \xi_k}{a_n} \Rightarrow S_\alpha(t), \quad t \in [0, 1],$$

where $S_\alpha(t)$ is a homogeneous vector-valued process in $D(\mathbb{R}^d)$ with independent increments and $S_\alpha(1)$ distributed according to a stable law of index α . Then (similarly to (3.11)), asymptotic equivalence (3.4) along with the suitable modification of Theorem 14.4 in [6, p. 152] implies

$$\frac{\sum_{k=0}^{c(\lfloor nt \rfloor)-1} \xi_k}{a_n} \Rightarrow b^{1/\alpha} \cdot S_\alpha(t), \quad t \in [0, 1],$$

where $b = \frac{\pi_1}{E(T_1)}$. In particular, using $t = 1$,

$$(3.16) \quad \frac{\sum_{k=0}^{c(n)-1} \xi_k}{a_n} \Rightarrow b^{1/\alpha} \cdot S_\alpha(1),$$

Recall r_k from (3.6). Since

$$\left\| X_t - \sum_{k=0}^{c(t)-1} \xi_k \right\| \leq r_{c(t)},$$

an application of the renewal theorem shows that

$$\frac{X_n}{n} \Rightarrow \mathcal{L}_\alpha \quad \text{and hence} \quad \frac{X_{\lfloor t \rfloor}}{t} \Rightarrow \mathcal{L}_\alpha.$$

Since $\|X_{\lfloor t \rfloor} - X_t\| \leq 1$, the proof of part (a)-(i) of Theorem 2.5 is completed. \square

Part (a)-(ii) of Theorem 2.5. For $v \in U$ let $c_v(t)$ be the number of occurrences of v in the set $\{\eta_1, \eta_2, \dots, \eta_{N_t}\}$. That is,

$$c_v(t) = \sum_{k=1}^{N_t} \mathbf{1}_{\{\eta_k=v\}}, \quad t \geq 0, v \in U.$$

Notice that $c_{u_1}(t) = c(t)$, where $c(t)$ is introduced in Section 3.1. Similarly to (3.4) we have

$$(3.17) \quad \lim_{t \rightarrow \infty} \frac{c_v(t)}{t} = \frac{\pi_v}{E(T_1)}, \quad \text{a. s.},$$

where π_v is the mass that the stationary distribution of the Markov chain η puts on v .

Define $\tau_v(0) = 0$ and $\tau_v(j) = \inf\{k > \tau_v(j-1) : \eta_k = v\}$ for $j \in \mathbb{N}$. For $v \in U$ and $t \geq 0$, let

$$\tilde{B}_v(t) = \sum_{i=0}^{c_v(t)-1} T_{\tau_v(i)} - c_v(t) \cdot E(T_1).$$

Then, the law of iterated logarithm for heavy-tailed i.i.d. sequences (see Theorems 1.6.6 and 3.9.1 in [7]) combined with (3.4) yields

$$(3.18) \quad \limsup_{t \rightarrow \infty} \frac{\tilde{B}_v(t)}{a_t \cdot (\ln t)^{1/\alpha + \varepsilon}} = \begin{cases} 0 & \text{if } \varepsilon > 0, \\ \infty & \text{if } \varepsilon < 0 \end{cases} \quad \text{a. s.}$$

For $v \in U$, let

$$B_v(t) = \sum_{i=0}^{c_v(t)-1} (T_{\tau_v(i)} - E(T_1)) + (t - s_{N_t}) \mathbf{1}_{\{\eta_{N_t} = v\}}.$$

Then, (1.1) implies that

$$X_t = \sum_{v \in U} v B_v(t) + \left(\sum_{v \in U} v \cdot c_v(t) \cdot E(T_1) - \mu \cdot t \right).$$

Taking into account (3.17), a standard inversion argument allows one to deduce from the law of iterated logarithm for $\tau_v(n)$ that

$$(3.19) \quad \limsup_{t \rightarrow \infty} \frac{\left\| \sum_{v \in U} v \cdot c_v(t) \cdot E(T_1) - \mu \cdot t \right\|}{\sqrt{t \ln \ln t}} < \infty, \quad \text{a. s.}$$

Since $a_t \in \mathcal{R}_\alpha$ with $\alpha \in (1, 2)$,

$$\lim_{t \rightarrow \infty} \frac{\sqrt{t \ln \ln t}}{a_t \cdot (\ln t)^{1/\alpha + \varepsilon}} = 0.$$

Thus (3.18) along with (3.19) yields part (a)-(ii) of Theorem 2.5, provided that we are able to show that for any $u, v \in U$ and all $\delta \in (1/(2\alpha), 1/\alpha)$,

$$(3.20) \quad P(E_{n,v,\delta} \cap E_{n,u,\delta} \text{ i. o.}) = 0,$$

where the events $E_{n,v,\delta}$ are defined for $n \in \mathbb{N}$, $v \in U$, and $\delta \in (1/(2\alpha), 1/\alpha)$ as follows:

$$E_{n,v,\delta} := \left\{ \max_{1 \leq m \leq c_v(n)} \left| \sum_{i=0}^{m-1} T_{\tau_v(i)} - m \cdot E(T_1) \right| > a_n \cdot (\ln n)^\delta \right\}.$$

For $n \in \mathbb{N}$ let $\gamma_n = 2n \cdot \max_{v \in V} \pi_v$ and define

$$G_{n,v,\delta} := \left\{ \max_{1 \leq m \leq \gamma_n} \left| \sum_{i=0}^{m-1} T_{\tau_v(i)} - m \cdot E(T_1) \right| > a_n \cdot (\ln n)^\delta \right\}.$$

Fix now any directions $u, v \in U$ and a constant $\delta \in (1/(2\alpha), 1/\alpha)$. We have:

$$\begin{aligned} & P(E_{n,v,\delta} \cap E_{n,u,\delta}) \\ & \leq P(G_{n,v,\delta} \cap G_{n,u,\delta}) + P(c_v(n) > \gamma_n) + P(c_u(n) > \gamma_n) \\ & = P(G_{n,v,\delta}) \cdot P(G_{n,u,\delta}) + P(c_v(n) > \gamma_n) + P(c_u(n) > \gamma_n). \end{aligned}$$

It follows from the large deviation principle for the random sequence $c_v(n)/n$ that $P(c_v(n) > \gamma_n) < K_v e^{-n\lambda_v}$ for some $K_v > 0$ and $\lambda_v > 0$. Furthermore, for any $A > 0$ and $k_n = \lfloor A^n \rfloor$, we have $P(G_{k_n,v,\delta}) \leq C n^{-\beta}$ for some constants $\beta = \beta(\delta) > 1/2$ and $C > 0$ (see [7, p. 177]; here we exploit the constraint $2\alpha\delta > 1$). The Borel-Cantelli lemma implies then that $P(G_{k_n,v,\delta} \cap G_{k_n,u,\delta} \text{ i. o.}) = 0$. Since for any $n \in \mathbb{N}$ there is a unique $j(n) \in \mathbb{N}$ such that $k_{j(n)} \leq n < k_{j(n)+1}$, and $\lim_{k \rightarrow \infty} \frac{a_{k+1} (\ln a_{k+1})^\delta}{a_k (\ln a_k)^\delta} = 1$, this yields (3.20). The proof of part (a)-(ii) of Theorem 2.5 is therefore completed. \square

3.5. Proof of Theorem 2.6. Define two sequences of processes in $D(\mathbb{R})$, $(B_n)_{n \in \mathbb{N}}$ and $(C_n)_{n \in \mathbb{N}}$, by setting

$$(3.21) \quad B_n(t) := \frac{\sum_{k=1}^{\lfloor nt \rfloor} \xi_k}{a_n} \quad \text{and} \quad C_n(t) := \frac{s_{\tau_{\lfloor nt \rfloor}}}{a_n}, \quad t \in [0, 1].$$

Lemma 3.1 combined with [4, Theorem 1.1] implies that $(\xi_1, s_{\tau_2} - s_{\tau_1}) \in \mathcal{R}_{d+1, \alpha, a}$, and hence

$$(3.22) \quad (B_n, C_n) \Rightarrow (S_\alpha, U_\alpha),$$

where S_α and U_α are homogeneous process with independent increments in $D(\mathbb{R}^d)$ and $D(\mathbb{R})$, respectively, such that $S_\alpha(1)$ and $U_\alpha(1)$ have (multivariate in the former case) stable distributions of index α .

Let U_n^{-1} and C_n^{-1} denote the inverse processes of U_n and C_n , respectively. One can define C_n^{-1} explicitly as follows:

$$(3.23) \quad C_n^{-1}(t) = n^{-1}c(a_nt), \quad t \in [0, 1].$$

Then the same argument as in [13, pp. 380-381] shows that (alternatively, one can use the result of [31]):

$$(B_n, C_n^{-1}) \Rightarrow (S_\alpha, U_\alpha^{-1})$$

in $D(\mathbb{R}^{d+1})$. This along with (3.23) implies (see, for instance, [6, p. 151]) that

$$\frac{\sum_{i=1}^{c(a_n)-1} \xi_i}{a_n} \Rightarrow \mathcal{L}_\alpha,$$

where

$$(3.24) \quad \mathcal{L}_\alpha := S_\alpha(U_\alpha^{-1})(1).$$

Passing to the subsequence $m_n = \lfloor a_n^{-1} \rfloor$ and using basic properties of regularly varying functions, we obtain

$$(3.25) \quad \frac{\sum_{i=0}^{c(n)-1} \xi_i}{n} \Rightarrow \mathcal{L}_\alpha.$$

To conclude the proof of the theorem one can use verbatim the argument along the lines following (3.16) in the concluding paragraph of the above proof of part (a)-(i) of Theorem 2.5. Namely, taking into account the inequality

$$\left\| X_t - \sum_{k=0}^{c(t)-1} \xi_k \right\| \leq r_{c(t)}$$

and using the renewal theorem which ensures the weak convergence of $r_{c(t)}$ to a proper random variable, (3.25) yields that $\frac{X_t}{t} \Rightarrow \mathcal{L}_\alpha$. The proof of Theorem 2.6 is completed. \square

ACKNOWLEDGEMENT

We would like to thank the anonymous Referee and the Associate Editor for their very helpful comments and suggestions.

REFERENCES

1. P. Allaart and M. Monticino, *Optimal stopping rules for directionally reinforced processes*, Adv. in Appl. Probab. **33** (2001), 483–504.
2. P. Allaart and M. Monticino, *Optimal buy/sell rules for correlated random walks*, J. Appl. Probab. **45** (2008), 33–44.
3. K. B. Athreya and P. E. Ney, *A new approach to the limit theory of recurrent Markov chains*, Trans. Amer. Math. Soc. **245** (1978), 493–501.
4. B. Basrak, R. A. Davis, and T. Mikosch, *A characterization of multivariate regular variation*, Ann. Appl. Probab. **12** (2000), 908–920.
5. P. Becker-Kern, M. M. Meerschaert, and H. P. Scheffler, *Limit theorems for coupled continuous time random walks*, Annals of Probab. **32** (2004), 730–756.
6. P. Billingsley, *Convergence of Probability Measures*, 2nd ed., (Wiley Series in Probability and Statistics), Wiley-Interscience, 1999.
7. A. A. Borovkov and K. A. Borovkov, *Asymptotic Analysis of Random Walks: Heavy-Tailed Distributions (Encyclopedia of Mathematics and its Applications 118)*, Cambridge University Press, 2008.
8. D. Buraczewski, E. Damek, and M. Mirek, *Asymptotics of stationary solutions of multivariate stochastic recursions with heavy tailed inputs and related limit theorems*, Stoch. Proc. Appl. **122** (2012), 42–67.
9. M. Csörgö and L. Horváth, *Weighted Approximations in Probability and Statistics*, Wiley, New York, 1993.
10. S. Goldstein, *On diffusion by discontinuous movements, and on the telegraph equation*, Quart. J. Mech. Appl. Math. **4** (1951), 129–156.
11. A. De Gregorio, *On random flights with non-uniformly distributed directions*, preprint is available at <http://arxiv.org/abs/1110.3621>.
12. U. Gruber and M. Schweizer, *A diffusion limit for generalized correlated random walks*, J. Appl. Probab. **43** (2006), 60–73.
13. L. Horváth and Q. M. Shao, *Limit distributions of directionally reinforced random walks*, Adv. Math. **134** (1998), 367–383.
14. A. Jurliewicz, P. Kern, M. M. Meerschaert, and H. P. Scheffler, *Fractional governing equations for coupled random walks*, Computers and Mathematics with Applications, to appear in the Special Issue on Advances in Fractional Differential Equations III (2012). Preprint available at <http://www.stt.msu.edu/~mcubed/OCTRW2.pdf>.
15. M. Kac, *A stochastic model related to the telegrapher's equation*, Rocky Mountain J. Math. **4** (1974), 497–509.
16. J. E. Kiefer and G. H. Weiss, *The Pearson random walk*, AIP Conf. Proc. **109** (1984), 11–32.
17. A. D. Kolesnik, *Random motion at finite speed in higher dimensions*, J. Stat. Phys. **131** (2008), 1039–1065.
18. A. D. Kolesnik and E. Orsingher, *A planar random motion with a infinite number of directions controlled by the damped wave equation*, J. Appl. Probab. **42** (2005), 1168–1182.
19. A. Lachal, *Cyclic random motions in space \mathbb{R}^d with n directions*, ESAIM Probab. Stat. **10** (2006), 277–316.
20. R. D. Mauldin, M. Monticino, and H. von Weizsäcker, *Directionally reinforced random walks*, Adv. Math. **117** (1996), 239–252.
21. M. M. Meerschaert and H. P. Scheffler, *Triangular array limits for continuous time random walks*, Stoch. Proc. Appl. **118** (2008), 1606–1633.
22. E. Orsingher and A. De Gregorio, *Random flights in higher spaces*, J. Theoret. Probab. **20** (2007), 769–806.
23. K. Pearson, *The problem of the random walk*, Nature **72** (1905), 294.
24. S. I. Resnick, *On the foundations of multivariate heavy tail analysis*, J. Appl. Probab. **41** (2004), 191–212.
25. R. Siegmund-Schultze and H. von Weizsäcker, *Level crossing probabilities II: Polygonal recurrence of multidimensional random walks*, Adv. Math. **208** (2007), 680–698.
26. W. Stadje, *The exact probability distribution of a two-dimensional random walk*, J. Stat. Phys. **46** (1987), 207–216.
27. P. Straka and B. I. Henry, *Lagging and leading coupled continuous time random walks, renewal times and their joint limits*, Stoch. Proc. Appl. **121** (2011), 324–336.

28. D. Szász and B. Tóth, *Persistent random walks in a one-dimensional random environment*, J. Stat. Phys. **37** (1984), 27–38.
29. B. Tóth, *Persistent random walks in random environment*, Probab. Theory Related Fields **71** (1986), 615–625.
30. B. J. West, *Statistics of deep water surface waves*. In *Encyclopedia of Fluid Mechanics, Vol. 2: Dynamics of Single Fluid Flows and Mixing* (N. P. Cheremisinoff, Ed.), Gulf, Houston, 1986, pp. 26–46.
31. W. Whitt, *Weak convergence of first passage time processes*, J. Appl. Probab. **8** (1971), 417–422.

DEPARTMENT OF STATISTICS, IOWA STATE UNIVERSITY, AMES, IA 50011, USA

E-mail address: `apghosh@iastate.edu`

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IA 50011, USA

E-mail address: `rastegar@iastate.edu`

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IA 50011, USA

E-mail address: `roiterst@iastate.edu`