

Discrete Structures for Computing

CSCE 222

Sandeep Kumar

Many slides based on [Lee19], [Rog21], [GK22]

Relations

Chapter 8

©2019 McGraw-Hill Education. All rights reserved. Authorized only for instructor use in the classroom. No reproduction or further distribution permitted without the prior written consent of McGraw-Hill Education.

Chapter Summary

- Relations and Their Properties
- n -ary Relations and Their Applications
- Representing Relations
- Closures of Relations
- Equivalence Relations
- Partial Orderings

Relations and Their Properties

Section 8.1


- Relations and Functions
- Properties of Relations
 - ▶ Reflexive Relations
 - ▶ Symmetric and Antisymmetric Relations
 - ▶ Transitive Relations
- Combining Relations

Binary Relations

Definition: A binary relation R from a set A to a set B is a subset $R \subseteq A \times B$.

Example:

- Let $A = \{0, 1, 2\}$ and $B = \{a, b\}$.
- $\{(0, a), (0, b), (1, a), (2, b)\}$ is a relation from A to B .
- We can represent relations from a set A to a set B graphically or using a table:



R	a	b
0	×	×
1	×	
2		×

Relations are more general than functions. A function is a relation where exactly one element of B is related to each element of A .

Binary Relations on a Set

Definition: A binary relation R on a set A is a subset of $A \times A$ or a relation from A to A .

Example:

- Suppose that $A = \{a, b, c\}$. Then $R = \{(a, a), (a, b), (a, c)\}$ is a relation on A .
- Let $A = \{1, 2, 3, 4\}$. The ordered pairs in the relation $R = \{(a, b) \mid a \text{ divides } b\}$ are

$(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3),$ and $(4, 4)$

Binary Relations on a Set

How many relations are there on a set A ?

Because a relation on A is the same thing as a subset of $A \times A$,

- We count the subsets of $A \times A$.
- Since $A \times A$ has n^2 elements when A has n elements, and
- A set with m elements has 2^m subsets,
- There are 2^{n^2} subsets of $A \times A$.
- Therefore, there are 2^{n^2} relations on a set A .

Binary Relations on a Set

Example: Consider these relations on the set of integers:

$$R_1 = \{(a, b) \mid a \leq b\},$$

$$R_2 = \{(a, b) \mid a > b\},$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a, b) \mid a = b\},$$

$$R_5 = \{(a, b) \mid a = b + 1\},$$

$$R_6 = \{(a, b) \mid a + b \leq 3\}.$$

Note that these relations are on an infinite set and each of these relations is an infinite set.

- Which of these relations contain each of the pairs

$(1, 1)$, $(1, 2)$, $(2, 1)$, $(1, -1)$, and $(2, 2)$?

- Checking the conditions that define each relation, we see that
 - ▶ the pair $(1, 1)$ is in R_1 , R_3 , R_4 , and R_6 ,
 - ▶ $(1, 2)$ is in R_1 and R_6 ,
 - ▶ $(2, 1)$ is in R_2 , R_5 , and R_6 ,
 - ▶ $(1, -1)$ is in R_2 , R_3 , and R_6 ,
 - ▶ $(2, 2)$ is in R_1 , R_3 , and R_4 .

Reflexive Relations

Definition: R is *reflexive* iff $(a, a) \in R$ for every element $a \in A$. Written symbolically, R is reflexive if and only if

$$\forall x [x \in A \rightarrow (x, x) \in R]$$

The following relations on the integers are reflexive:

$$R_1 = \{(a, b) \mid a \leq b\},$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a, b) \mid a = b\}$$

The following relations are *not* reflexive:

$$R_2 = \{(a, b) \mid a > b\} \quad (\text{note that } 3 \not> 3)$$

$$R_5 = \{(a, b) \mid a = b + 1\} \quad (\text{note that } 3 \neq 3 + 1)$$

$$R_6 = \{(a, b) \mid a + b \leq 3\} \quad (\text{note that } 4 + 4 \not\leq 3)$$

If $A = \emptyset$ then the empty relation is reflexive vacuously. That is, the empty relation on an empty set is reflexive!

Symmetric Relations

Definition: R is symmetric iff $(b, a) \in R$ whenever $(a, b) \in R$ for all $(a, b) \in A$. Written symbolically, R is symmetric if and only if

$$\forall x \forall y [(x, y) \in R \rightarrow (y, x) \in R]$$

The following relations on the integers are symmetric:

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a, b) \mid a = b\},$$

$$R_6 = \{(a, b) \mid a + b \leq 3\}.$$

The following are *not* symmetric:

$$R_1 = \{(a, b) \mid a \leq b\} \quad (\text{note that } 3 \leq 4, \text{ but } 4 \not\leq 3),$$

$$R_2 = \{(a, b) \mid a > b\} \quad (\text{note that } 4 > 3, \text{ but } 3 \not> 4),$$

$$R_5 = \{(a, b) \mid a = b + 1\} \quad (\text{note that } 4 = 3 + 1, \text{ but } 3 \neq 4 + 1)$$

Antisymmetric Relations

Definition: A relation R on a set A such that for all $a, b \in A$ if $(a, b) \in R$ and $(b, a) \in R$, then $a = b$ is called *antisymmetric*. Written symbolically, R is antisymmetric if and only if

$$\forall x \forall y [(x, y) \in R \wedge (y, x) \in R \rightarrow x = y]$$

The following relations on the integers are antisymmetric.

$$R_1 = \{(a, b) \mid a \leq b\},$$

$$R_2 = \{(a, b) \mid a > b\},$$

$$R_4 = \{(a, b) \mid a = b\},$$

$$R_5 = \{(a, b) \mid a = b + 1\}.$$

The following relations are *not* antisymmetric.

- $R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$. Note that both $(1, -1)$ and $(-1, 1)$ belong to R_3 .
- $R_6 = \{(a, b) \mid a + b \leq 3\}$. Note that both $(1, 2)$ and $(2, 1)$ belong to R_6 .

Transitive Relations

Definition: A relation R on a set A is called transitive if whenever $(a, b) \in R$ and $(b, c) \in R$, then $(a, c) \in R$, for all $a, b, c \in A$. Written symbolically, R is transitive if and only if

$$\forall x \forall y \forall z [(x, y) \in R \wedge (y, z) \in R \rightarrow (x, z) \in R]$$

The following relations on the integers are transitive. Note that for every integer, $a \leq b$ and $b \leq c$, then $a \leq c$.

$$R_1 = \{(a, b) \mid a \leq b\},$$

$$R_2 = \{(a, b) \mid a > b\},$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a, b) \mid a = b\}.$$

The following are not transitive.

- $R_5 = \{(a, b) \mid a = b + 1\}$. Note that both $(3, 2)$ and $(4, 3)$ belongs to R_5 , but not $(4, 2)$.
- $R_6 = \{(a, b) \mid a + b \leq 3\}$. Note that both $(2, 1)$ and $(1, 2)$ belongs to R_6 , but not $(2, 2)$.

Combining Relations

Given two relations R_1 and R_2 , we can combine them using basic set operations to form new relations such as $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, and $R_2 - R_1$.

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 3, 4\}$. The relations

- $R_1 = \{(1, 1), (2, 2), (3, 3)\}$, and
- $R_2 = \{(1, 1), (1, 2), (1, 3), (1, 4)\}$

can be combined using basic set operations to form new relations:

$$R_1 \cup R_2 = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (3, 3)\}$$

$$R_1 \cap R_2 = \{(1, 1)\}$$

$$R_1 - R_2 = \{(2, 2), (3, 3)\}$$

$$R_2 - R_1 = \{(1, 2), (1, 3), (1, 4)\}$$

Composition


Definition: Suppose

- R_1 is a relation from a set A to a set B .
- R_2 is a relation from B to a set C .

Then the *composition* (or *composite*) of R_2 with R_1 , is a relation from A to C where

- if (x, y) is a member of R_1 and (y, z) is a member of R_2 , then (x, z) is a member of $R_2 \circ R_1$.

Representing the Composition of Relations



$$R_2 \circ R_1 = \{(b, x), (b, z)\}$$

Powers of a Relation

Definition: Let R be a binary relation on A . Then the powers R^n of the relation R can be defined inductively by:

- Basis Step: $R^1 = R$.
- Inductive Step: $R^{n+1} = R^n \circ R$

The powers of a transitive relation are subsets of the relation. This is established by the following theorem:

Theorem 1: The relation R on a set A is transitive iff $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

See the text for a proof via mathematical induction.

Representing Relations

Section 8.3 Summary

- Representing Relations using Matrices
- Representing Relations using Digraphs

Representing Relations Using Matrices

- A relation between finite sets can be represented using a zero-one matrix.
- Suppose R is a relation from $A = \{a_1, a_2, \dots, a_m\}$ to $B = \{b_1, b_2, \dots, b_n\}$.
 - ▶ The elements of the two sets can be listed in any particular arbitrary order. When $A = B$, we use the same ordering.
- The relation R is represented by the matrix $M_R = [m_{ij}]$, where

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R, \\ 0 & \text{if } (a_i, b_j) \notin R. \end{cases}$$

- The matrix representing R has a 1 as its (i, j) entry when a_i is related to b_j and a 0 if a_i is not related to b_j .

Examples of Representing Relations Using Matrices

Suppose that $A = \{1, 2, 3\}$ and $B = \{1, 2\}$. Let R be the relation from A to B containing (a, b) if $a \in A$, $b \in B$, and $a > b$. What is the matrix representing R assuming the ordering of elements is in increasing numerical order?

Because $R = \{(2, 1), (3, 1), (3, 2)\}$, the matrix is

$$M_R = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}$$

Examples of Representing Relations Using Matrices

Example: Let $A = \{a_1, a_2, a_3\}$ and $B = \{b_1, b_2, b_3, b_4, b_5\}$. Which ordered pairs are in the relation R represented by the matrix

$$M_R = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix}$$

Because R consists of those ordered pairs (a_i, b_j) with $m_{ij} = 1$, it follows that:

$$R = \{(a_1, b_2), (a_2, b_1), (a_2, b_3), (a_2, b_4), (a_3, b_1), (a_3, b_3), (a_3, b_5)\}$$



Matrices of Relations on Sets

If R is a reflexive relation, all the elements on the main diagonal of M_R are equal to 1.

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & \ddots & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

R is a symmetric relation, if and only if $m_{ij} = 1$ whenever $m_{ji} = 1$.

R is an antisymmetric relation, if and only if $m_{ij} = 0$ or $m_{ji} = 0$ when $i \neq j$.



Example of a Relation on a Set

Suppose that the relation R on a set is represented by the matrix

$$M_R = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

Is R reflexive, symmetric, and/or antisymmetric?


- Because all the diagonal elements are equal to 1, R is reflexive.
- Because M_R is symmetric, R is symmetric and,
- It's not antisymmetric because both $m_{1,2}$ and $m_{2,1}$ are 1.

Representing Relations Using Digraphs

Definition: A *directed graph*, or *digraph*, consists of a set V of vertices (or nodes) together with a set E of ordered pairs of elements of V called *edges* (or arcs).


- The vertex a is called the initial vertex of the edge (a, b) , and
- The vertex b is called the terminal vertex of this edge.
- An edge of the form (a, a) is called a loop.

Example: A drawing of the directed graph with vertices a, b, c , and d , and edges (a, b) , (a, d) , (b, b) , (b, d) , (c, a) , (c, b) , and (d, b) is shown here.



Examples of Digraphs Representing Relations

Example: What are the ordered pairs in the relation represented by this directed graph?




The ordered pairs in the relation are:

- $(1, 3)$, $(1, 4)$,
- $(2, 1)$, $(2, 2)$, $(2, 3)$,
- $(3, 1)$, $(3, 3)$,
- $(4, 1)$, and $(4, 3)$.

Determining which Properties a Relation has from its Digraph


- *Reflexivity*: A loop must be present at all vertices in the graph.
- *Symmetry*: If (x, y) is an edge, then so is (y, x) .
- *Antisymmetry*: If (x, y) with $x \neq y$ is an edge, then (y, x) is not an edge.
- *Transitivity*: If (x, y) and (y, z) are edges, then so is (x, z) .

Determining which Properties a Relation has from its Digraph




- Reflexive?
 - ▶ No, not every vertex has a loop
- Symmetric?
 - ▶ Yes (trivially), there is no edge from one vertex to another.
- Antisymmetric?
 - ▶ Yes (trivially), there is no edge from one vertex to another.
- Transitive?
 - ▶ Yes, (trivially) since there is no edge from one vertex to another.

Determining which Properties a Relation has from its Digraph







- Reflexive?
 - ▶ No, there are no loops.
- Symmetric?
 - ▶ No, there is an edge from a to b , but not from b to a .
- Antisymmetric?
 - ▶ No, there is an edge from d to b and b to d .
- Transitive?
 - ▶ No, there are edges from a to b and from b to d , but there is no edge from a to d .

Determining which Properties a Relation has from its Digraph



- Reflexive?
 - ▶ No, there are no loops
- Symmetric?
 - ▶ No, for example, there is no edge from c to a .
- Antisymmetric?
 - ▶ Yes, whenever there is an edge from one vertex to another, there is not one going back.
- Transitive?
 - ▶ Yes.

Example of the Powers of a Relation



The pair (x, y) is in R^n if there is a path of length n from x to y in R , following the direction of the arrows.

Section Summary

Section 8.5

- Equivalence Relations
- Equivalence Classes
- Equivalence Classes and Partitions

Equivalence Relations

Definition 1: A relation on a set A is called an *equivalence relation* if it is reflexive, symmetric, and transitive.

Definition 2: Two elements a and b that are related by an equivalence relation are called *equivalent*. The notation $a \sim b$ is often used to denote that a and b are equivalent elements with respect to a particular equivalence relation.

Strings

Suppose that R is the relation on the set of strings of English letters such that aRb if and only if $\ell(a) = \ell(b)$, where $\ell(x)$ is the length of the string x . Is R an equivalence relation?

Show that all of the properties of an equivalence relation hold.

- *Reflexivity*: Because $\ell(a) = \ell(a)$, it follows that aRa for all strings a .
- *Symmetry*: Suppose that aRb . Since $\ell(a) = \ell(b)$, $\ell(b) = \ell(a)$ also holds and bRa .
- *Transitivity*: Suppose that aRb and bRc . Since $\ell(a) = \ell(b)$, and $\ell(b) = \ell(c)$, $\ell(a) = \ell(c)$ also holds and aRc .

Congruence Modulo m

Let m be an integer > 1 . Show that the relation

$$R = \{(a, b) \mid a \equiv b \pmod{m}\}$$

is an equivalence relation on the set of integers.

Recall that $a \equiv b \pmod{m}$ if and only if m divides $a - b$.

- *Reflexivity*: $a \equiv a \pmod{m}$ since $a - a = 0$ is divisible by m since $0 = 0 \cdot m$.
- *Symmetry*: Suppose that $a \equiv b \pmod{m}$. Then $a - b$ is divisible by m , and so $a - b = km$, where k is an integer. It follows that $b - a = (-k)m$, so $b \equiv a \pmod{m}$.
- *Transitivity*: Suppose that $a \equiv b \pmod{m}$ and $b \equiv c \pmod{m}$. Then m divides both $a - b$ and $b - c$. Hence, there are integers k and ℓ with $a - b = km$ and $b - c = \ell m$. We obtain by adding the equations:

$$a - c = (a - b) + (b - c) = km + \ell m = (k + \ell)m$$

- Therefore, $a \equiv c \pmod{m}$.

Divides

Show that the *divides* relation on the set of positive integers is *not* an equivalence relation.

The properties of reflexivity, and transitivity do hold, but the relation is not symmetric. Hence, divides is not an equivalence relation.

- *Reflexivity*: $a \mid a$ for all a .
- *Not Symmetric*: For example, $2 \mid 4$, but $4 \nmid 2$. Hence, the relation is not symmetric.
- *Transitivity*: Suppose that a divides b and b divides c . Then there are positive integers k and ℓ such that $b = ak$ and $c = b\ell$. Hence, $c = a(k\ell)$, so a divides c . Therefore, the relation is transitive.

Equivalence Classes

Definition 3: Let R be an equivalence relation on a set A . The set of all elements that are related to an element a of A is called the equivalence class of a . The equivalence class of a with respect to R is denoted by $[a]_R$.

When only one relation is under consideration, we can write $[a]$, without the subscript R , for this equivalence class. Note that

$$[a]_R = \{s \mid (a, s) \in R\}$$

If $b \in [a]_R$, then b is called a representative of this equivalence class. Any element of a class can be used as a representative of the class.

The equivalence classes of the relation congruence modulo m are called the *congruence classes modulo m* . The congruence class of an integer a modulo m is denoted by $[a]_m$, so

$$[a]_m = \{\dots, a - 2m, a - m, a, a + m, a + 2m, \dots\}$$

For example,

- $[0]_4 = \{\dots, -8, -4, 0, 4, 8, \dots\}$
- $[1]_4 = \{\dots, -7, -3, 1, 5, 9, \dots\}$

Equivalence Classes and Partitions

Theorem 1: Let R be an equivalence relation on a set A . These statements for elements a and b of A are equivalent:

- 1 aRb
- 2 $[a] = [b]$
- 3 $[a] \cap [b] \neq \phi$

Proof: We show that (i) implies (ii).


- Assume that aRb .
- Now suppose that $c \in [a]$. Then aRc .
- Because aRb and R is symmetric, bRa .
- Because R is transitive and bRa and aRc , it follows that bRc .
- Hence, $c \in [b]$. Therefore, $[a] \subseteq [b]$.
- A similar argument shows that $[b] \subseteq [a]$.
- Since $[a] \subseteq [b]$ and $[b] \subseteq [a]$, we have shown that $[a] = [b]$.

See text for proof that (ii) implies (iii) and (iii) implies (i).

Partition of a Set

Definition: A *partition* of a set S is a collection of disjoint nonempty subsets of S that have S as their union. In other words, the collection of subsets A_i , where $i \in I$ (where I is an index set), forms a partition of S if and only if

- $A_i \neq \phi$ for $i \in I$,
- $A_i \cap A_j = \phi$ when $i \neq j$,
- and $\cup_{i \in I} A_i = S$.



An Equivalence Relation Partitions a Set

Let R be an equivalence relation on a set A . The union of all the equivalence classes of R is all of A , since an element a of A is in its own equivalence class $[a]_R$. In other words,

$$\bigcup_{a \in A} [a]_R = A$$

From Theorem 1, it follows that these equivalence classes are either equal or disjoint, so

$$[a]_R \cap [b]_R = \phi \text{ when } [a]_R \neq [b]_R$$

Therefore, the equivalence classes form a partition of A , because they split A into disjoint subsets.

An Equivalence Relation Partitions a Set

Theorem 2: Let R be an equivalence relation on a set S . Then the equivalence classes of R form a partition of S .

Conversely, given a partition $\{A_i \mid i \in I\}$ of the set S , there is an equivalence relation R that has the sets $A_i, i \in I$, as its equivalence classes.

Proof: We have already shown the first part of the theorem.

For the second part, assume that $\{A_i \mid i \in I\}$ is a partition of S . Let R be the relation on S consisting of the pairs (x, y) where x and y belong to the same subset A_i in the partition. We must show that R satisfies the properties of an equivalence relation.

An Equivalence Relation Partitions a Set

- *Reflexivity*: For every $a \in S$, $(a, a) \in R$, because a is in the same subset as itself.
- *Symmetry*: If $(a, b) \in R$, then b and a are in the same subset of the partition, so $(b, a) \in R$.
- *Transitivity*: If $(a, b) \in R$ and $(b, c) \in R$, then a and b are in the same subset of the partition, as are b and c .

Since the subsets are disjoint and b belongs to both, the two subsets of the partition must be identical.

Therefore, $(a, c) \in R$ since a and c belong to the same subset of the partition.

Partial Orderings

Section 8.6

- Partial Orderings and Partially-ordered Sets
- Lexicographic Orderings
- Hasse Diagrams
- Lattices
- Topological Sorting

Partial Orderings

Definition 1: A relation R on a set S is called a *partial ordering*, or *partial order*, if it is

- reflexive,
- antisymmetric, and
- transitive.

A set together with a partial ordering R is called a *partially ordered set*, or *poset*, and is denoted (S, R) . Members of S are called *elements* of the poset.

Partial Orderings

Example 1: Show that the *greater than or equal* relation (\geq) is a partial ordering on the set of integers.

- *Reflexivity:* $a \geq a$ for every integer a .
- *Antisymmetry:* If $a \geq b$ and $b \geq a$, then $a = b$.
- *Transitivity:* If $a \geq b$ and $b \geq c$, then $a \geq c$.


These properties all follow from the order axioms for the integers. (See Appendix 1).

Partial Orderings

Example 2: Show that the divisibility relation ($|$) is a partial ordering on the set of integers.

- *Reflexivity:* $a | a$ for all integers a . (See Example 9 in Section 8.1).
- *Antisymmetry:* If a and b are positive integers with $a | b$ and $b | a$, then $a = b$. (See Example 12 in Section 9.1)
- *Transitivity:* Suppose that a divides b and b divides c . Then there are positive integers k and ℓ such that $b = ak$ and $c = b\ell$. Hence, $c = a(k\ell)$, so a divides c . Therefore, the relation is transitive.

$(\mathbb{Z}^+, |)$ is a poset.




Partial Orderings

Example 3: Show that the inclusion relation (\subseteq) is a partial ordering on the power set of a set S .

- *Reflexivity:* $A \subseteq A$ whenever A is a subset of S .
- *Antisymmetry:* If A and B are positive integers with $A \subseteq B$ and $B \subseteq A$, then $A = B$.
- *Transitivity:* If $A \subseteq B$ and $B \subseteq C$, then $A \subseteq C$.

The properties all follow from the definition of set inclusion.



Comparability

Definition 2: The elements a and b of a poset (S, \preceq) are *comparable* if either $a \preceq b$ or $b \preceq a$. When a and b are elements of S so that neither $a \preceq b$ nor $b \preceq a$, then a and b are called *incomparable*.

The symbol \preceq is used to denote the relation in any poset.

Definition 3: If (S, \preceq) is a poset and every two elements of S are comparable, S is called a *totally ordered* or *linearly ordered* set, and is called a *total order* or a *linear order*. A totally ordered set is also called a *chain*.

Definition 4: (S, \preceq) is a well-ordered set if it is a poset such that \preceq is a total ordering and every nonempty subset of S has a least element.

- Does total ordering imply well ordering?

Lexicographic Order

Definition: Given two posets (A_1, \preceq_1) and (A_2, \preceq_2) , the *lexicographic ordering* on $A_1 \times A_2$ is defined by specifying that:

- (a_1, a_2) is less than (b_1, b_2) , i.e., $(a_1, a_2) \prec (b_1, b_2)$, either if
 - ▶ $a_1 \prec_1 b_1$ or
 - ▶ $a_1 = b_1$ and $a_2 \prec_2 b_2$.

Bard: How do you construct a partial ordering from the cross product of two posets?

Example: Consider strings of lowercase English letters. A lexicographic ordering can be defined using the ordering of the letters in the alphabet. This is the same ordering that is used in dictionaries.

- *discreet* \prec *discrete*, because these strings differ in the seventh position and $e \prec t$.
- *discreet* \prec *discreetness*, because the first eight letters agree, but the second string is longer.


Hasse Diagrams

Definition: A *Hasse diagram* is a visual representation of a partial ordering that leaves out edges that must be present because of the reflexive and transitive properties.

A partial ordering is shown in (a) of the figure (next slide).

- The loops due to the reflexivity property are deleted in (b).
- The edges that must be present due to the transitive property are deleted in (c).
- The Hasse diagram for the partial ordering (a), is depicted in (c).

Hasse Diagrams



Procedure for Constructing a Hasse Diagram

To represent a finite poset (S, \preceq) using a Hasse diagram, start with the directed graph of the relation:

- Remove the loops (a, a) present at every vertex due to the reflexive property.
- Remove all edges (x, y) for which there is an element $z \in S$ such that $x \prec z$ and $z \prec y$. These are the edges that must be present due to the transitive property.
- Arrange each edge so that its initial vertex is below the terminal vertex. Remove all the arrows, because all edges point upwards toward their terminal vertex.

Bibliography I



Ashutosh Gupta and S. Krishna.

Cs 228: Logic for computer science 2022.

<https://www.cse.iitb.ac.in/~akg/courses/2022-logic/>, January 2022.



Hyunyoung Lee.

Discrete structures for computing.

Class slides for TAMU CSCE 222, 2019.



Phillip Rogaway.

Ecs20 fall 2021 lecture notes, Fall 2021.