

# THE “FULL CLARKSON-ERDŐS-SCHWARTZ THEOREM” ON THE CLOSURE OF NON-DENSE MÜNTZ SPACES

TAMÁS ERDÉLYI

ABSTRACT. Denote by  $\text{span}\{f_1, f_2, \dots\}$  the collection of all finite linear combinations of the functions  $f_1, f_2, \dots$  over  $\mathbb{R}$ . The principal result of the paper is the following.

**Theorem (Full Clarkson-Erdős-Schwartz Theorem).** *Suppose  $(\lambda_j)_{j=1}^{\infty}$  is a sequence of distinct positive numbers. Then  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  is dense in  $C[0, 1]$  if and only if*

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} = \infty.$$

Moreover, if

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} < \infty,$$

then every function from the  $C[0, 1]$  closure of  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  can be represented as an analytic function on  $\{z \in \mathbb{C} \setminus (-\infty, 0] : |z| < 1\}$  restricted to  $(0, 1)$ .

This result improves an earlier result by P. Borwein and Erdélyi stating that if

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} < \infty,$$

then every function from the  $C[0, 1]$  closure of  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  is in  $C^{\infty}(0, 1)$ . Our result may also be viewed as an improvement, extension, or completion of earlier results by Müntz, Szász, Clarkson, Erdős, L. Schwartz, P. Borwein, Erdélyi, W.B. Johnson, and Operstein.

---

1991 *Mathematics Subject Classification*. Primary: 30B60, 41A17.

*Key words and phrases*. Müntz Theorem, denseness in  $C[0, 1]$ , Erdos-Clarkson-Schwartz Theorem.

Research of T. Erdélyi is supported, in part, by NSF under Grant No. DMS-0070826.

## 1. INTRODUCTION AND NOTATION

Müntz's beautiful classical theorem characterizes sequences  $(\lambda_j)_{j=0}^\infty$  with

$$0 = \lambda_0 < \lambda_1 < \lambda_2 < \dots$$

for which the Müntz space  $\text{span}\{x^{\lambda_0}, x^{\lambda_1}, \dots\}$  is dense in  $C[0, 1]$ . Here, and in what follows,  $\text{span}\{x^{\lambda_0}, x^{\lambda_1}, \dots\}$  denotes the collection of finite linear combinations of the functions  $x^{\lambda_0}, x^{\lambda_1}, \dots$  with real coefficients, and  $C[a, b]$  is the space of all real-valued continuous functions on  $[a, b] \subset \mathbb{R}$  equipped with the uniform norm. Müntz's Theorem [Bo-Er3, De-Lo, Go, Mü, Szá] states the following.

**Theorem 1.A (Müntz).** *Suppose  $(\lambda_j)_{j=0}^\infty$  is a sequence with  $0 = \lambda_0 < \lambda_1 < \lambda_2 < \dots$ . Then  $\text{span}\{x^{\lambda_0}, x^{\lambda_1}, \dots\}$  is dense in  $C[0, 1]$  if and only if  $\sum_{j=1}^\infty 1/\lambda_j = \infty$ .*

The original Müntz Theorem proved by Müntz [Mü] in 1914, by Szász [Szá] in 1916, and anticipated by Bernstein [Be] was only for sequences of exponents tending to infinity. The point 0 is special in the study of Müntz spaces. Even replacing  $[0, 1]$  by an interval  $[a, b] \subset [0, \infty)$  in Müntz's Theorem is a non-trivial issue. This is, in large measure, due to Clarkson and Erdős [Cl-Er] and Schwartz [Sch] whose works include the result that if  $\sum_{j=1}^\infty 1/\lambda_j < \infty$  then every function belonging to the uniform closure of  $\text{span}\{x^{\lambda_0}, x^{\lambda_1}, \dots\}$  on  $[a, b]$  can be extended analytically throughout the region  $\{z \in \mathbb{C} \setminus (-\infty, 0] : |z| < b\}$ .

There are many variations and generalizations of Müntz's Theorem [An, Be, Boa, Bo1, Bo2, Bo-Er1, Bo-Er2, Bo-Er3, Bo-Er4, Bo-Er5, Bo-Er6, Bo-Er7, B-E-Z, Ch, Cl-Er, De-Lo, Er-Jo, Go, Lu-Ko, Ma, Op, Sch, So]. There are also still many open problems. In [Bo-Er6] it is shown that the interval  $[0, 1]$  in Müntz's Theorem can be replaced by an arbitrary compact set  $A \subset [0, \infty)$  of positive Lebesgue measure. That is, if  $A \subset [0, \infty)$  is a compact set of positive Lebesgue measure, then  $\text{span}\{x^{\lambda_0}, x^{\lambda_1}, \dots\}$  is dense in  $C(A)$  if and only if  $\sum_{j=1}^\infty 1/\lambda_j = \infty$ . Here  $C(A)$  denotes the space of all real-valued continuous functions on  $A$  equipped with the uniform norm. If  $A$  contains an interval then this follows from the already mentioned results of Clarkson, Erdős, and Schwartz. However, their results and methods cannot handle the case when, for example,  $A \subset [0, 1]$  is a Cantor type set of positive measure.

In the case that  $\sum_{j=1}^\infty 1/\lambda_j < \infty$ , analyticity properties of the functions belonging to the uniform closure of  $\text{span}\{x^{\lambda_0}, x^{\lambda_1}, \dots\}$  on  $A$  are also established in [Bo-Er6].

In [Bo-Er3, Section 4.2] and in [Bo-Er4] the following result is proved.

**Theorem 1.B (Full Müntz Theorem in  $C[0, 1]$ ).** *Suppose  $(\lambda_j)_{j=1}^\infty$  is a sequence of distinct positive real numbers. Then  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  is dense in  $C[0, 1]$  if and only if*

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} = \infty.$$

Moreover, if

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} < \infty,$$

then every function from the  $C[0, 1]$  closure of  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  is infinitely many times differentiable on  $(0, 1)$ .

The new result of this paper is the following.

**Theorem 1.1 (Full Clarkson-Erdős-Schwartz Theorem).** *Suppose  $(\lambda_j)_{j=1}^{\infty}$  is a sequence of distinct positive numbers. Then  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  is dense in  $C[0, 1]$  if and only if*

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} = \infty.$$

Moreover, if

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} < \infty,$$

then every function from the  $C[0, 1]$  closure of  $\text{span}\{1, x^{\lambda_1}, x^{\lambda_2}, \dots\}$  can be represented as an analytic function on  $\{z \in \mathbb{C} \setminus (-\infty, 0] : |z| < 1\}$  restricted to  $(0, 1)$ .

The notation

$$\|f\|_A := \sup_{x \in A} |f(x)|$$

is used throughout this paper for real-valued measurable functions  $f$  defined on a set  $A \subset \mathbb{R}$ . The space of all real-valued continuous functions on a set  $A \subset \mathbb{R}$  equipped with the uniform norm is denoted by  $C(A)$ . Denote by  $\text{span}\{f_1, f_2, \dots\}$  the collection of all finite linear combinations of the functions  $f_1, f_2, \dots$  over  $\mathbb{R}$ .

## 2. AUXILIARY RESULTS

The following result is the “bounded Remez-type inequality for non-dense Müntz spaces” due to P. Borwein and Erdélyi [Bo-Er6].

**Theorem 2.1.** *Suppose  $(\gamma_j)_{j=1}^{\infty}$  is a sequence of distinct positive numbers satisfying*

$$\sum_{j=1}^{\infty} 1/\gamma_j < \infty.$$

Let  $s > 0$ . Then there exists a constant  $c(\Gamma, s)$  depending only on  $\Gamma := (\gamma_j)_{j=1}^{\infty}$  and  $s$  (and not on  $\varrho$ ,  $A$ , or the “length” of  $Q$ ) so that

$$\|Q\|_{[0, \varrho]} \leq c(\Gamma, s) \|Q\|_A$$

for every  $Q \in \text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\}$  and for every set  $A \subset [\varrho, 1]$  of Lebesgue measure at least  $s$ .

Combining a result of Clarkson and Erdős [Cl-Er] and its extension given by Schwartz [Sch] we can state the following

**Theorem 2.2.** *Suppose  $(\gamma_j)_{j=1}^{\infty}$  is a sequence of distinct positive numbers satisfying  $\sum_{j=1}^{\infty} 1/\gamma_j < \infty$ . Then  $\text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\}$  is not dense in  $C[0, 1]$ . In addition, if the gap condition*

$$\inf\{\gamma_{j+1} - \gamma_j : j = 1, 2, \dots\} > 0$$

*holds, then every function  $f \in C[0, 1]$  belonging to the  $C[0, 1]$  closure of  $\text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\}$  can be represented as*

$$f(x) = \sum_{j=0}^{\infty} a_j x^{\gamma_j}, \quad x \in [0, 1].$$

*If the gap condition (2.1) does not hold, then every function  $f \in C[0, 1]$  belonging to the  $C[0, 1]$  closure of  $\text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\}$  can still be represented as an analytic function on*

$$\{z \in \mathbb{C} \setminus (-\infty, 0] : |z| < 1\}$$

*restricted to  $(0, 1)$ .*

Now we offer a sufficient condition for a sequence  $(\beta_j)_{j=1}^{\infty}$  of distinct positive numbers converging to 0 to guarantee the non-denseness of  $\text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}$  in  $C[0, 1]$ .

**Theorem 2.3.** *Suppose that  $(\beta_j)_{j=1}^{\infty}$  is a sequence of distinct real numbers greater than 0 satisfying*

$$\sum_{j=1}^{\infty} \beta_j =: \eta < \infty.$$

*Then  $\text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}$  is not dense in  $C[0, 1]$ . In addition, every function in the  $C[0, 1]$  closure of  $\text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}$  can be represented as an analytic function on  $\mathbb{C} \setminus (-\infty, 0]$  restricted to  $(0, 1)$ .*

*Proof of Theorem 2.3.* The theorem is a consequence of D. J. Newman's Markov-type inequality [Bo-Er3, Theorem 6.1.1 on page 276] (see also [Ne]). We state this as Theorem 2.4. Repeated applications of Theorem 2.4 with the substitution  $x = e^{-t}$  imply that

$$\|Q(e^{-t})^{(m)}\|_{[0, \infty)} \leq (9\eta)^m \|Q(e^{-t})\|_{[0, \infty)}, \quad m = 1, 2, \dots,$$

in particular

$$|(Q(e^{-t})^{(m)})(0)| \leq (9\eta)^m \|Q(e^{-t})\|_{[0, \infty)}, \quad m = 1, 2, \dots,$$

for every  $Q \in \text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}$ . By using the Taylor series expansion of  $Q(e^{-t})$  around 0, we obtain that

$$(2.1) \quad |Q(z)| \leq c_1(K, \eta) \|Q\|_{[0, 1]}, \quad z \in K,$$

for every  $Q \in \text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}$  and for every compact  $K \subset \mathbb{C} \setminus \{0\}$ , where

$$c_1(K, \eta) := \sum_{m=0}^{\infty} \frac{(9\eta)^m \left( \max_{z \in K} |\log z| \right)^m}{m!} = \exp \left( 9\eta \max_{z \in K} |\log z| \right)$$

is a constant depending only on  $K$  and  $\eta$ . Now (2.1) shows that if

$$Q_n \in \text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}$$

converges in  $C[0, 1]$ , then it converges uniformly on every compact  $K \subset \mathbb{C} \setminus \{0\}$ , and the theorem is proved.  $\square$

The following Markov-type inequality for Müntz polynomials is due to Newman [Bo-Er3, Theorem 6.1.1 on page 276] (see also [Ne]).

**Theorem 2.4 (Markov-Type Inequality for Müntz Polynomials).** *Suppose that  $\beta_1, \beta_2, \dots, \beta_n$  are distinct nonnegative numbers. Then*

$$\|xQ'(x)\|_{[0,1]} \leq 9 \left( \sum_{j=1}^n \beta_j \right) \|Q\|_{[0,1]}$$

for every  $Q \in \text{span}\{x^{\beta_1}, x^{\beta_2}, \dots, x^{\beta_n}\}$ .

We will also need the bounded Bernstein-type inequality below (see [Bo-Er3, page 182]).

**Theorem 2.5 (Bernstein Type Inequality for Non-Dense Müntz spaces).** *Suppose  $\Gamma := (\gamma_j)_{j=1}^{\infty}$  is a sequence of distinct positive numbers satisfying  $\gamma_1 \geq 1$  and  $\sum_{j=1}^{\infty} 1/\gamma_j < \infty$ . Then*

$$\|Q'\|_{[0,x]} \leq c(\Gamma, x) \|Q\|_{[0,1]}$$

for every  $Q \in \text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\}$  and for every  $x \in [0, 1)$ , where  $c(\Gamma, x)$  depends only on  $\Gamma$  and  $x$ .

The following simple fact will also be needed.

**Lemma 2.6.** *Let  $U \subset C[0, 1]$  be a closed linear subspace and let  $V \subset C[0, 1]$  be a finite dimensional (hence closed) linear subspace. Then  $U + V$  is closed.*

#### 4. PROOF OF THEOREMS 1.1

*Proof of Theorem 1.1.* The first part of the theorem is contained in Theorem 1.B, so we need to prove only the second part. Suppose  $(\lambda_j)_{j=1}^{\infty}$  is a sequence of distinct positive numbers satisfying

$$\sum_{j=1}^{\infty} \frac{\lambda_j}{\lambda_j^2 + 1} < \infty.$$

Then there are positive numbers  $\eta$ ,  $\beta_j$ ,  $\gamma_j$ , and  $\delta_j$  such that

$$\{\lambda_j : j = 1, 2, \dots\} = \{\beta_j : j = 1, 2, \dots\} \cup \{\gamma_j : j = 1, 2, \dots\} \cup \{\delta_j : j = 1, 2, \dots, k\},$$

where  $\gamma_1 \geq 1$ ,

$$\sum_{j=1}^{\infty} \beta_j \leq \eta, \quad \sum_{j=1}^{\infty} 1/\gamma_j < \infty,$$

and with  $\Gamma := (\gamma_j)_{j=1}^{\infty}$  we have

$$c(\Gamma, 1/2) < \frac{1}{36\eta}$$

( $c(\Gamma, 1/2)$  is defined in Theorem 2.1). Let

$$H_{\beta} := \text{span}\{x^{\beta_1}, x^{\beta_2}, \dots\}, \quad H_{\gamma} := \text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\},$$

and

$$H_{\delta} := \text{span}\{x^{\delta_1}, x^{\delta_2}, \dots, x^{\delta_k}\}.$$

Every  $Q \in H_{\beta} + H_{\gamma}$  can be written as  $Q = Q_{\beta} + Q_{\gamma}$  with some  $Q_{\beta} \in H_{\beta}$  and  $Q_{\gamma} \in H_{\gamma}$ . First we show that there are constants  $C_{\beta}$  and  $C_{\gamma}$  depending only on  $H_{\beta}$  and  $H_{\gamma}$ , respectively, so that

$$(3.1) \quad \|Q_{\beta}\|_{[0,1]} \leq C_{\beta} \|Q\|_{[0,1]}$$

and

$$(3.2) \quad \|Q_{\gamma}\|_{[0,1]} \leq C_{\gamma} \|Q\|_{[0,1]}$$

for every  $Q \in H_{\beta} + H_{\gamma}$ . Suppose to the contrary that, say the first inequality fails. Then there are Müntz polynomials  $Q_{\beta,n} \in H_{\beta}$  and  $Q_{\gamma,n} \in H_{\gamma}$  so that

$$(3.3) \quad \|Q_{\beta,n}\|_{[0,1]} = 1, \quad \lim_{n \rightarrow \infty} \|Q_{\gamma,n}\|_{[0,1]} = 1,$$

and

$$(3.4) \quad \lim_{n \rightarrow \infty} \|Q_{\beta,n} + Q_{\gamma,n}\|_{[0,1]} = 0.$$

Then by Theorem 2.4  $\{Q_{\beta,n} : n = 1, 2, \dots\}$  is a family of bounded, equi-continuous functions on  $[1/3, 1]$ , while  $\{Q_{\gamma,n} : n = 1, 2, \dots\}$  is a family of bounded, equi-continuous functions on  $[0, 2/3]$ . So by the Arzela-Ascoli Theorem there are a subsequence of  $(Q_{\beta,n})$  (without loss of generality we may assume that this is  $(Q_{\beta,n})$  itself) and a subsequence of  $(Q_{\gamma,n})$  (without loss of generality we may assume that this is  $(Q_{\gamma,n})$  itself) so that

$$(3.5) \quad \lim_{n \rightarrow \infty} \|Q_{\beta,n} - f\|_{[1/3,1]} = 0$$

and

$$(3.6) \quad \lim_{n \rightarrow \infty} \|Q_{\gamma,n} - g\|_{[0,2/3]} = 0$$

with some continuous functions  $f$  and  $g$  on  $[1/3, 1]$  and  $[0, 2/3]$ , respectively. By (3.4), (3.5), and (3.6) we have  $f = -g$  on  $[1/3, 2/3]$ , so the function

$$(3.7) \quad h(x) := \begin{cases} f(x), & x \in [1/3, 1] \\ -g(x), & x \in [0, 2/3] \end{cases}$$

is well-defined. By (3.4) – (3.7) we can deduce that

$$(3.8) \quad \lim_{n \rightarrow \infty} \|Q_{\beta, n} - h\|_{[0,1]} = 0$$

and

$$(3.9) \quad \lim_{n \rightarrow \infty} \|Q_{\gamma, n} - h\|_{[0,1]} = 0.$$

Using (3.3), (3.8), Theorem 2.4, and  $\sum_{j=1}^{\infty} \beta_j \leq \eta$  we can deduce that

$$h(x) - h(1) \leq 18\eta, \quad x \in [1/2, 1].$$

Note that (3.3), (3.5), and (3.7) imply that  $\|h\|_{[0,1]} = 1$  and  $h(0) = 0$ . Now observe that the function  $h - h(1)$  is in the uniform closure of

$$H_{\gamma} = \text{span}\{1, x^{\gamma_1}, x^{\gamma_2}, \dots\},$$

hence Theorem 2.1 implies

$$\|h - h(1)\|_{[0,1]} \leq c(\Gamma, 1/2) \|h - h(1)\|_{[1/2,1]} \leq c(\Gamma, 1/2) 18\eta < 1/2.$$

This contradicts the facts that  $h(0) = 0$  and  $\|h\|_{[0,1]} = 1$ . Hence the proof of (3.1) is finished. The proof of (3.2) goes in the same way, so we omit it.

Let  $\overline{H}$  denote the uniform closure of a subspace  $H \subset C[0, 1]$ . We want to prove that  $\overline{H_{\beta} + H_{\gamma} + H_{\delta}} \subset \mathcal{A}$ , where  $\mathcal{A} \subset C[0, 1]$  denotes the collection of functions  $f \in C[0, 1]$ , which can be represented as an analytic function on  $\{z \in \mathbb{C} \setminus (-\infty, 0] : |z| < 1\}$  restricted to  $(0, 1)$ . Since  $H_{\delta}$  is finite dimensional, Theorem 2.6 implies that

$$\overline{H_{\beta} + H_{\gamma} + H_{\delta}} \subset \overline{H_{\beta} + H_{\gamma}} + H_{\delta}.$$

so it is sufficient to prove that

$$(3.10) \quad \overline{H_{\beta} + H_{\gamma}} \subset \mathcal{A}$$

However, (3.1) and (3.2) imply that

$$\overline{H_{\beta} + H_{\gamma}} \subset \overline{H_{\beta}} + \overline{H_{\gamma}},$$

where  $\overline{H_{\beta}} \subset \mathcal{A}$  by Theorem 2.3 and  $\overline{H_{\gamma}} \subset \mathcal{A}$  by Theorem 2.2. Hence (3.10) holds, indeed, and the proof of the theorem is finished.  $\square$

## REFERENCES

- [An] J.M. Anderson, *Müntz-Szász type approximation and the angular growth of lacunary integral functions*, Trans. Amer. Math. Soc. **169** (1972), 237–248.
- [Ba] J. Bastero,  *$\ell_q$ -subspaces of stable  $p$ -Banach spaces*, Archiv. Math. **40** (1983), 538–544.
- [Be] S.N. Bernstein, *Collected Works: Vol 1. Constructive Theory of Functions (1905-1930), English Translation*, Atomic Energy Commission, Springfield, Va, 1958.
- [Boa] R.P. Boas, *Entire Functions*, Academic Press, New York, 1954.
- [Bo1] P.B. Borwein, *Zeros of Chebyshev polynomials in Markov Systems*, J. Approx. Theory **63** (1990), 56–64.
- [Bo2] P.B. Borwein, *Variations on Müntz’s theme*, Can. Math. Bull. **34** (1991), 305–310.
- [Bo-Er1] P.B. Borwein & T. Erdélyi, *Notes on lacunary Müntz polynomials* **76** (1991), Israel J. Math., 183–192.
- [Bo-Er2] P.B. Borwein & T. Erdélyi, *Lacunary Müntz systems*, Proc. Edinburgh Math. Soc. **36** (1993), 361–374.
- [Bo-Er3] P.B. Borwein & T. Erdélyi, *Polynomials and Polynomial Inequalities*, Springer-Verlag, Graduate Texts in Mathematics, 486 p., 1995.
- [Bo-Er4] P.B. Borwein & T. Erdélyi, *The full Müntz theorem in  $C[0, 1]$  and  $L_1[0, 1]$* , J. London Math. Soc. **54** (1996), 102–110.
- [Bo-Er5] P.B. Borwein & T. Erdélyi, *The  $L_p$  version of Newman’s inequality for lacunary polynomials*, Proc. Amer. Math. Soc. **124** (1996), 101–109.
- [Bo-Er6] P.B. Borwein & T. Erdélyi, *Generalizations of Müntz’s theorem via a Remez-type inequality for Müntz spaces*, J. Amer. Math. Soc. **10** (1997), 327–329.
- [Bo-Er7] P.B. Borwein & T. Erdélyi, *Müntz’s Theorem on compact subsets of positive measure*, in Approximation Theory, eds. N.K. Govil et al., Marcel Dekker, Inc. (1998), 115–131.
- [B-E-Z] P.B. Borwein, T. Erdélyi, & J. Zhang, *Müntz systems and orthogonal Müntz-Legendre polynomials* **342** (1994), Trans. Amer. Math. Soc., 523–542.
- [Ch] E.W. Cheney, *Introduction to Approximation Theory*, McGraw-Hill, New York, 1966.
- [Cl-Er] J.A. Clarkson & P. Erdős, *Approximation by polynomials*, Duke Math. J. **10** (1943), 5–11.
- [De-Lo] R.A. DeVore & G.G. Lorentz, *Constructive Approximation*, Springer-Verlag, Berlin, 1993.
- [Er-Jo] T. Erdélyi & W.B. Johnson, *The “Full Müntz Theorem” in  $L_p[0, 1]$  for  $0 < p < \infty$* , Journal D’Analyse Mathématique **84** (2001), 145–172.
- [Go] M. von Golitschek, *A short proof of Müntz Theorem*, J. Approx. Theory **39** (1983), 394–395.
- [Lu-Ko] W.A.J. Luxemburg & J. Korevaar, *Entire functions and Müntz-Szász type approximation*, Trans. Amer. Math. Soc. **157** (1971), 23–37.
- [Ma] S. Mazur, *On the convergence of lacunary polynomials*, Studia Math. **89** (1988), 75–78.
- [Mü] C. Müntz, *Über den Approximationsatz von Weierstrass*, H.A. Schwartz Festschrift, Berlin (1914).
- [Ne] D.J. Newman, *Derivative bounds for Müntz polynomials*, J. Approx. Theory **18** (1976), 360–362.
- [Op] V. Operstein, *Full Müntz theorem in  $L_p[0, 1]$* , J. Approx. Theory **85** (1996), 233–235.



- [Ro] H.L. Royden, *Real Analysis, third ed.*, Macmillan, New York, NY, 1988.
- [Sch] L. Schwartz, *Etude des Sommes d'Exponentielles*, Hermann, Paris, 1959.
- [So] G. Somorjai, *A Müntz-type problem for rational approximation*, Acta. Math. Hungar. **27** (1976), 197–199.
- [Szá] O. Szász, *Über die Approximation stetiger Funktionen durch lineare Aggregate von Potenzen*, Math. Ann. **77** (1916), 482–496.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843  
*E-mail address:* `terdelyi@math.tamu.edu` (Tamás Erdélyi)