

A Sharp Remez Inequality on the Size of Constrained Polynomials

TAMÁS ERDÉLYI

*Department of Mathematics, The Ohio State University,
Columbus, Ohio 43210-1174, U.S.A.*

Communicated by V. Totik

Received June 30, 1989

Denote by Π_n the set of all real algebraic polynomials of degree at most n . We define the class

$$\Pi_n(s) = \{p \in \Pi_n : m(\{x \in [-1, 1] : |p(x)| \leq 1\}) \geq 2 - s\} \quad (0 < s < 2),$$

where $m(A)$ denotes the Lebesgue measure of A . How large can the maximum modulus be on $[-1, 1]$ for polynomials from $\Pi_n(s)$? In [7] E. J. Remez answered this question establishing the best possible upper bound. The solution and one of its applications in the theory of orthogonal polynomials can be found in [5] as well. Remez-type inequalities and their applications were studied in [1-3]. The purpose of this paper is to prove a sharp Remez-type inequality for constrained polynomials.

Remez's inequality asserts that

$$\max_{-1 \leq x \leq 1} |p(x)| \leq Q_n(4/(2-s) - 1) \quad (p \in \Pi_n(s), 0 < s < 2), \quad (1)$$

where $Q_n(x) = \cos(n \arccos x)$ is the Chebyshev polynomial of degree n . For $a < b$ we define

$$P_n(a, b) = \left\{ p : p(x) = \sum_{j=0}^n \alpha_j (b-x)^j (x-a)^{n-j} \text{ with all } \alpha_j \geq 0 \text{ or all } \alpha_j \leq 0 \right\}.$$

The class $P_n(-1, 1)$ was introduced and examined thoroughly by G. G. Lorentz in [6], subsequently a number of properties were obtained in [4]. By an observation of Lorentz, if $p \in \Pi_n$ has no zero in the open unit circle then $p \in P_n(-1, 1)$. In this paper we prove the following sharp Remez-type theorem for polynomials from $P_n(-1, 1)$.

THEOREM. *We have*

$$\max_{-1 \leq x \leq 1} |p(x)| \leq (1 - s/2)^{-n} \quad (p \in P_n(-1, 1) \cap \Pi_n(s), 0 < s < 2), \quad (2)$$

and the equality holds only for the polynomials $\pm(1 \pm x)^n/(2 - s)^n$.

COROLLARY. *If $p \in \Pi_n(s)$ has no zero in the open unit circle then (2) holds.*

Proof of the Theorem. Observe that $[c, d] \subset [a, b]$ implies $P_n(a, b) \subset P_n(c, d)$. This follows simply from the definition and the substitutions

$$\begin{aligned} b - x &= \frac{b - c}{d - c} (d - x) + \frac{b - d}{d - c} (x - c), \\ x - a &= \frac{c - a}{d - c} (d - x) + \frac{d - a}{d - c} (x - c), \end{aligned}$$

where $(b - c)/(d - c)$, $(b - d)/(d - c)$, $(c - a)/(d - c)$, and $(d - a)/(d - c)$ are non-negative. Let $p \in P_n(a, b)$ with the representation

$$p(x) = \sum_{j=0}^n \alpha_j (b - x)^j (x - a)^{n-j} \quad \text{with all } \alpha_j \geq 0 \text{ or all } \alpha_j \leq 0. \quad (3)$$

Then for $0 < s < 2$ we easily deduce

$$\begin{aligned} |p(b)| &= |\alpha_0| (b - a)^n = \left(\frac{b - a}{y - a} \right)^n |\alpha_0| (y - a)^n \leq \left(\frac{b - a}{y - a} \right)^n |p(y)| \\ &\leq (1 - s/2)^{-n} |p(y)| \quad (b - (b - a)s/2 \leq y \leq b) \end{aligned} \quad (4)$$

and similarly

$$|p(a)| \leq (1 - s/2)^{-n} |p(y)| \quad (a \leq y \leq a + (b - a)s/2). \quad (5)$$

Now let $p \in P_n(-1, 1) \cap \Pi_n(s)$ ($0 < s < 2$), and choose a $z \in [-1, 1]$ such that

$$|p(z)| = \max_{-1 \leq x \leq 1} |p(x)|. \quad (6)$$

Since $p \in \Pi_n(s)$, there is a y from either $[z - s(z + 1)/2, z]$ or $[z, z + s(1 - z)/2]$ such that $|p(y)| \leq 1$. In the first case the relation $P_n(-1, 1) \subset P_n(-1, z)$ and (4) yield the desired result, and in the second case the relation $P_n(-1, 1) \subset P_n(z, 1)$ and (5) give the theorem.

REFERENCES

1. T. ERDÉLYI, Remez-type inequality on the size of generalized polynomials, manuscript.
2. T. ERDÉLYI, The Remez inequality on the size of polynomials, in "Approximation Theory VI: I," C. K. Chui, L. L. Schumaker and J. D. Ward (Eds.), pp. 243–246.
3. T. ERDÉLYI, Nikolskii-type inequalities for generalized polynomials and zeros of orthogonal polynomials, *J. Approx. Theory*, to appear.
4. T. ERDÉLYI AND J. SZABADOS, On polynomials with positive coefficients, *J. Approx. Theory* **54** (1988), 107–122.
5. G. FREUD, "Orthogonal Polynomials," Pergamon, Oxford, 1971.
6. G. G. LORENTZ, The degree of approximation by polynomials with positive coefficients, *Math. Ann.* **151** (1963), 239–251.
7. E. J. REMEZ, Sur une propriété des polynômes de Tchebycheff, *Comm. Inst. Sci. Kharkow* **13** (1936), 93–95.