

# Coppersmith-Rivlin type inequalities and the order of vanishing of polynomials at 1

Tamás Erdélyi  
Department of Mathematics  
Texas A&M University  
College Station, Texas 77843

December 4, 2015

## Abstract

For  $n \in \mathbb{N}$ ,  $L > 0$ , and  $p \geq 1$  let  $\kappa_p(n, L)$  be the largest possible value of  $k$  for which there is a polynomial  $P \not\equiv 0$  of the form

$$P(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| \geq L \left( \sum_{j=1}^n |a_j|^p \right)^{1/p}, \quad a_j \in \mathbb{C},$$

such that  $(x-1)^k$  divides  $P(x)$ . For  $n \in \mathbb{N}$ ,  $L > 0$ , and  $q \geq 1$  let  $\mu_q(n, L)$  be the smallest value of  $k$  for which there is a polynomial  $Q$  of degree  $k$  with complex coefficients such that

$$|Q(0)| > \frac{1}{L} \left( \sum_{j=1}^n |Q(j)|^q \right)^{1/q}.$$

We find the size of  $\kappa_p(n, L)$  and  $\mu_q(n, L)$  for all  $n \in \mathbb{N}$ ,  $L > 0$ , and  $1 \leq p, q \leq \infty$ . The result about  $\mu_\infty(n, L)$  is due to Coppersmith and Rivlin, but our proof is completely different and much shorter even in that special case.

---

2010 *Mathematics Subject Classification*: 11C08, 41A17, 26C10, 30C15.

*Key words and phrases*: polynomials, restricted coefficients, order of vanishing at 1, Coppersmith-Rivlin type inequalities

# 1 Introduction.

In [B-99] and [B-13] we examined a number of problems concerning polynomials with coefficients restricted in various ways. We are particularly interested in how small such polynomials can be on the interval  $[0, 1]$ . For example, we proved that there are absolute constants  $c_1 > 0$  and  $c_2 > 0$  such that

$$\exp(-c_1\sqrt{n}) \leq \min_{0 \neq Q \in \mathcal{F}_n} \left\{ \max_{x \in [0,1]} |Q(x)| \right\} \leq \exp(-c_2\sqrt{n})$$

for every  $n \geq 2$ , where  $\mathcal{F}_n$  denotes the set of all polynomials of degree at most  $n$  with coefficients from  $\{-1, 0, 1\}$ .

Littlewood considered minimization problems of this variety on the unit disk. His most famous, now solved, conjecture was that the  $L_1$  norm of an element  $f \in \mathcal{F}_n$  on the unit circle grows at least as fast as  $c \log N$ , where  $N$  is the number of non-zero coefficients in  $f$  and  $c > 0$  is an absolute constant.

When the coefficients are required to be integers, the questions have a Diophantine nature and have been studied from a variety of points of view. See [A-90], [B-98], [B-95], [F-80], [O-93].

One key to the analysis is a study of the related problem of giving an upper bound for the multiplicity of the zero these restricted polynomials can have at 1. In [B-99] and [B-13] we answer this latter question precisely for the class of polynomials of the form

$$Q(x) = \sum_{j=0}^n a_j x^j, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}, \quad j = 1, 2, \dots, n,$$

with fixed  $|a_0| \neq 0$ .

Variants of these questions have attracted considerable study, though rarely have precise answers been possible to give. See in particular [A-79], [B-32], [B-87], [E-50], [Sch-33], [Sz-34]. Indeed, the classical, much studied, and presumably very difficult problem of Prouhet, Tarry, and Escott rephrases as a question of this variety. (Precisely: what is the maximal vanishing at 1 of a polynomial with integer coefficients with  $l_1$  norm  $2n$ ? It is conjectured to be  $n$ .) See [H-82], [B-94], or [B-02].

For  $n \in \mathbb{N}$ ,  $L > 0$ , and  $p \geq 1$  we define the following numbers. Let  $\kappa_p(n, L)$  be the largest possible value of  $k$  for which there is a polynomial  $Q \not\equiv 0$  of the form

$$Q(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| \geq L \left( \sum_{j=1}^n |a_j|^p \right)^{1/p}, \quad a_j \in \mathbb{C},$$

such that  $(x-1)^k$  divides  $Q(x)$ . For  $n \in \mathbb{N}$  and  $L > 0$  let  $\kappa_\infty(n, L)$  be the largest possible value of  $k$  for which there is a polynomial  $Q \not\equiv 0$  of the form

$$Q(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| \geq L \max_{1 \leq j \leq n} |a_j|, \quad a_j \in \mathbb{C},$$

such that  $(x-1)^k$  divides  $Q(x)$ . In [B-99] we proved that there is an absolute constant  $c_3 > 0$  such that

$$\min \left\{ \frac{1}{6} \sqrt{n(1 - \log L)} - 1, n \right\} \leq \kappa_\infty(n, L) \leq \min \left\{ c_3 \sqrt{n(1 - \log L)}, n \right\}$$

for every  $n \in \mathbb{N}$  and  $L \in (0, 1]$ . However, we were far from being able to establish the right result in the case of  $L \geq 1$ . In [B-13] we proved the right order of magnitude of  $\kappa_\infty(n, L)$  and  $\kappa_2(n, L)$  in the case of  $L \geq 1$ . Our results in [B-99] and [B-13] sharpen and generalize results of Schur [Sch-33], Amoroso [A-90], Bombieri and Vaaler [B-87], and Hua [H-82] who gave versions of this result for polynomials with integer coefficients. Our results in [B-99] have turned out to be related to a number of recent papers from a wide range of research areas. See [A-02], [B-98], [B-95], [B-96], [B-97a], [B-97b], [B-97], [B-00], [B-07], [B-08a], [B-13], [B-08b], [B-92], [B-94], [Bu-99], [C-02], [C-13], [C-10], [D-99], [D-01], [D-13], [D-03], [E-08a], [E-08b], [E-13], [F-00], [G-05], [K-04], [M-68], [M-03], [N-94], [O-93], [P-99], [P-13], [R-04], [R-07], [S-99], [T-07], [T-84], for example. More results on the zeros of polynomials with Littlewood-type coefficient constraints may be found in [E-02b]. Markov and Bernstein type inequalities under Erdős type coefficient constraints are surveyed in [E-02a].

For  $n \in \mathbb{N}$ ,  $L > 0$ , and  $q \geq 1$  we define the following numbers. Let  $\mu_q(n, L)$  be the smallest value of  $k$  for which there is a polynomial of degree  $k$  with complex coefficients such that

$$|Q(0)| > \frac{1}{L} \left( \sum_{j=1}^n |Q(j)|^q \right)^{1/q}.$$

For  $n \in \mathbb{N}$  and  $L > 0$  let  $\mu_\infty(n, L)$  be the smallest value of  $k$  for which there is a polynomial of degree  $k$  with complex coefficients such that

$$|Q(0)| > \frac{1}{L} \max_{j \in \{1, 2, \dots, n\}} |Q(j)|.$$

It is a simple consequence of Hölder's inequality (see Lemma 3.6) that

$$\kappa_p(n, L) \leq \mu_q(n, L)$$

whenever  $n \in \mathbb{N}$ ,  $L > 0$ ,  $1 \leq p, q \leq \infty$ , and  $1/p + 1/q = 1$ .

In this paper we find the size of  $\kappa_p(n, L)$  and  $\mu_q(n, L)$  for all  $n \in \mathbb{N}$ ,  $L > 0$ , and  $1 \leq p, q \leq \infty$ . The result about  $\mu_\infty(n, L)$  is due to Copper-Smith and Rivlin [C-92], but our proof presented in this paper is completely different and much shorter even in that special case. Our results in [B-99] may be viewed as finding the size of  $\kappa_\infty(n, L)$  and  $\mu_1(n, L)$  for all  $n \in \mathbb{N}$  and  $L \in (0, 1]$ . Our results in [B-13] may be viewed as finding the size of  $\kappa_\infty(n, L)$ ,  $\mu_1(n, L)$ ,  $\kappa_2(n, L)$ , and  $\mu_2(n, L)$  for all  $n \in \mathbb{N}$  and  $L > 0$ .

## 2 New Results.

We extend some of our main results in [B-13] to the case  $L \geq 1$ . Our main result is the following.

**Theorem 2.1.** *Let  $p \in (1, \infty]$  and  $q \in [1, \infty)$  satisfy  $1/p + 1/q = 1$ . There are absolute constants  $c_1 > 0$  and  $c_2 > 0$  such that*

$$\sqrt{n}(c_1 L)^{-q/2} - 1 \leq \kappa_p(n, L) \leq \mu_q(n, L) \leq \sqrt{n}(c_2 L)^{-q/2} + 2$$

for every  $n \in \mathbb{N}$  and  $L > 1/2$ , and

$$\begin{aligned} c_3 \min \left\{ \sqrt{n(-\log L)}, n \right\} &\leq \kappa_p(n, L) \leq \mu_q(n, L) \\ &\leq c_4 \min \left\{ \sqrt{n(-\log L)}, n \right\} + 4 \end{aligned}$$

for every  $n \in \mathbb{N}$  and  $L \in (0, 1/2]$ . Here  $c_1 := 1/53$ ,  $c_2 := 40$ ,  $c_3 := 2/7$ , and  $c_4 := 13$  are appropriate choices

**Theorem 2.2.** *There are absolute constants  $c_1 > 0$  and  $c_2 > 0$  such that*

$$c_1 \sqrt{n(1-L)} - 1 \leq \kappa_1(n, L) \leq \mu_\infty(n, L) \leq c_2 \sqrt{n(1-L)} + 1$$

for every  $n \in \mathbb{N}$  and  $L \in (1/2, 1]$ , and

$$\begin{aligned} c_3 \min \left\{ \sqrt{n(-\log L)}, n \right\} &\leq \kappa_1(n, L) \leq \mu_\infty(n, L) \\ &\leq c_4 \min \left\{ \sqrt{n(-\log L)}, n \right\} + 4 \end{aligned}$$

for every  $n \in \mathbb{N}$  and  $L \in (0, 1/2]$ . Note that  $\kappa_1(n, L) = \mu_\infty(n, L) = 0$  for every  $n \in \mathbb{N}$  and  $L > 1$ . Here  $c_1 := 1/5$ ,  $c_2 := 1$ ,  $c_3 := 2/7$ , and  $c_4 := 13$  are appropriate choices.

### 3 Lemmas.

In this section we list our lemmas needed in the proofs of Theorems 2.1 and 2.2. These lemmas are proved in Section 4. Let  $\mathcal{P}_n$  be the set of all polynomials of degree at most  $n$  with real coefficients. Let  $\mathcal{P}_n^c$  be the set of all polynomials of degree at most  $n$  with complex coefficients.

**Lemma 3.1.** *Let  $p \in (1, \infty)$ . For any  $1 \leq M$  there are polynomials  $P_n$  of the form*

$$P_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad a_{0,n} \geq \frac{3M}{\pi^2} + o(M),$$

$$\left( \sum_{j=1}^n |a_{j,n}|^p \right)^{1/p} \leq 16M^{1/p},$$

such that  $P_n$  has at least  $\lfloor \sqrt{n/M} \rfloor$  zeros at 1.

**Lemma 3.2.** *Let  $p, q \in (1, \infty)$  satisfy  $1/p + 1/q = 1$ . For any  $L \geq 1/48$  there are polynomials  $P_n$  of the form*

$$P_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad a_{0,n} \geq L + o(L), \quad \sum_{j=1}^n |a_{j,n}|^p \leq 1,$$

such that  $P_n$  has at least  $\lfloor \sqrt{n}(cL)^{-q/2} \rfloor$  zeros at 1 with  $c := \frac{3}{16\pi^2}$ .

**Lemma 3.3.** *Let  $p \in [1, \infty)$ . For any  $L \in (0, 1/17)$  there are polynomials  $P_n$  of the form*

$$P_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad a_{0,n} = L, \quad \sum_{j=1}^n |a_{j,n}|^p \leq 1,$$

such that  $P_n$  has at least  $\frac{2}{7} \min\{\sqrt{n(1 - \log L)}, n\}$  zeros at 1.

**Lemma 3.4.** *For any  $L \in (0, 1)$  there are polynomials  $P_n \not\equiv 0$  of the form*

$$P_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad a_{0,n} \geq L \sum_{j=1}^n |a_{j,n}|,$$

such that  $P_n$  has at least  $\frac{1}{5} \sqrt{(n-1)(1-L)}$  zeros at 1.

The observation below is well known, easy to prove, and recorded in several papers. See [B-99], for example.

**Lemma 3.5.** *Let  $P \neq 0$  be a polynomial of the form  $P(x) = \sum_{j=0}^n a_j x^j$ . Then  $(x-1)^k$  divides  $P$  if and only if  $\sum_{j=0}^n a_j Q(j) = 0$  for all polynomials  $Q \in \mathcal{P}_{k-1}^c$ .*

Our next lemma is a simple consequence of Hölder's inequality.

**Lemma 3.6.** *Let  $1 \leq p, q \leq \infty$  and  $1/p + 1/q = 1$ . Then for every  $n \in \mathbb{N}$  and  $L > 0$ , we have*

$$\kappa_p(n, L) \leq \mu_q(n, L).$$

The next lemma is stated as Lemma 3.4 in [K-03], where a proof of it is also presented.

**Lemma 3.7.** *Given  $A > 0$  and  $M > 0$ , there exists a polynomial  $G$  such that  $F = G^2 \in \mathcal{P}_m$  with*

$$m < \sqrt{\pi} \sqrt{A} \sqrt[4]{M} + 2$$

such that  $F(0) = M$  and

$$|F(x)| \leq \min\{M, x^{-2}\}, \quad x \in (0, A].$$

We also need Lemma 5.7 from [B-99] which may be stated as follows.

**Lemma 3.8.** *Let  $n$  and  $R$  be positive integers with  $1 \leq R \leq \sqrt{n}$ . Then there exists a polynomial  $F \in \mathcal{P}_m$  with*

$$m \leq 4\sqrt{n} + \frac{9}{7}R\sqrt{n} + R + 4 \leq \frac{44}{7}R\sqrt{n} + 4$$

such that

$$F(1) = F(2) = \dots = F(R^2) = 0$$

and

$$\begin{aligned} |F(0)| &> \exp(R^2) (|F(R^2+1)| + |F(R^2+2)| + \dots + |F(n)|) \\ &\geq \exp(R^2) \left( \sum_{j=1}^n |F(j)|^2 \right)^{1/2}. \end{aligned}$$

Lemmas 3.6 and 3.7 imply the following results needed in the proof of Theorems 2.1 and 2.2.

**Lemma 3.9.** *Let  $q \in [1, \infty)$ . For every  $n \in \mathbb{N}$ ,  $q \in [1, \infty)$ , and  $K > 0$ , there are polynomials  $F \in \mathcal{P}_m$  satisfying*

$$|F(0)| > K \left( \sum_{j=1}^n |F(j)|^q \right)^{1/q}$$

and

$$m \leq \begin{cases} \sqrt{n}(40K)^{q/2} + 2, & 0 < K < 2, \\ 13 \min \{ \sqrt{n \log K}, n \} + 4, & K \geq 2. \end{cases}$$

**Lemma 3.10.** *For every  $n \in \mathbb{N}$  and  $K > 1$ , there are polynomials  $F \in \mathcal{P}_m$  satisfying*

$$|F(0)| > K \max_{j \in \{1, 2, \dots, n\}} |F(j)|$$

and

$$m \leq \begin{cases} \sqrt{n(K-1)/2} + 1, & 1 < K < 2, \\ 13 \min \{ \sqrt{n \log K}, n \} + 4, & K \geq 2. \end{cases}$$

## 4 Proofs of the Lemmas.

### Proof of Lemma 3.1.

Modifying the construction on page 138 of [B-95] for  $x \in (0, \infty)$  we define  $H_1(x) := 1$  and

$$H_m(x) := \frac{(-1)^{m+1} 2(m!)^2}{2\pi i} \int_{\Gamma} \frac{x^t dt}{(t-2) \prod_{j=0}^m (t-j^2)}, \quad m = 2, 3, \dots,$$

where the simple closed contour  $\Gamma$  surrounds the zeros of the denominator of the integrand. Observe that we have added the factor  $t-2$  to the denominator and the factor  $(-1)^{m+1} 2$  to achieve our goals. (It is left to the reader to see what role this modification plays in our proof.) Then  $H_m$  is a polynomial of degree  $m^2$  with a zero at 1 with multiplicity at least  $m+1$ . (This can be seen easily by repeated differentiation and then evaluation of the above contour integral by expanding the contour to infinity.)

Also, by the residue theorem,

$$(4.1) \quad H_m(x) = 1 + d_m x^2 + \sum_{k=1}^m c_{k,m} x^{k^2}, \quad m = 2, 3, \dots,$$

where

$$c_{k,m} = \frac{(-1)^{m+1} 2(m!)^2}{(k^2-2) \prod_{j=0, j \neq k}^m (k^2-j^2)} = \frac{4}{k^2-2} \frac{(-1)^{k+1} (m!)^2}{(m-k)!(m+k)!},$$

and

$$d_m = \frac{(-1)^{m+1} 2(m!)^2}{\prod_{j=0}^m (2-j^2)}.$$

It follows that each  $c_{k,m}$  is real and

$$(4.2) \quad |c_{k,m}| \leq \frac{4}{|k^2 - 2|}, \quad k = 1, 2, \dots, m,$$

and a simple calculation shows that

$$(4.3) \quad |d_m| \leq 8, \quad m = 2, 3, \dots$$

(No effort has been made to optimize the bound in (4.3).) Let  $S_M$  be the collection of all odd square free integers in  $[1, M]$ . Let  $m := \lfloor \sqrt{n/M} \rfloor$ . If  $m = 0$  then there is nothing to prove. So we may assume that  $m \geq 1$ . It is well known that

$$|S_M| \geq \frac{3M}{\pi^2} + o(M),$$

where  $|A|$  denotes the number of elements in a finite set  $A$ . This follows from the fact that if  $S_M^*$  is the collection of all square free integers in  $[1, M]$ , then

$$|S_M^*| = \frac{6M}{\pi^2} + o(M),$$

see pp. 267-268 in [H-38], for example, by observing that the number of odd square free integer in  $[1, M]$  is not less than the number of even square free integers in  $[1, M]$  (if  $a$  is an even square free integer then  $a/2$  is an odd square free integer). We define

$$P_n(x) := \sum_{j \in S_M} H_m(x^j).$$

Then  $P_n$  is of the form

$$P_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad j = 0, 1, \dots, n.$$

We have

$$a_{0,n} = |S_M^*| \geq \frac{3M}{\pi^2} + o(M).$$

First assume that  $m = 1$ . Then

$$\sum_{j=1}^n |a_{j,n}|^p = 2|S_M| \leq 2M,$$



and as  $P_n$  has 1 zero at 1, the lemma follows. Now assume that  $m \geq 2$ . We have  $ju \neq lv$  whenever  $j, l \in S_M$ ,  $j \neq l$ , and  $u, v \in \{1^2, 2^2, \dots, m^2\} \cup \{2\}$ . Combining this with (4.1), (4.2), and (4.3), we obtain

$$\begin{aligned} \sum_{j=1}^n |a_{j,n}|^p &\leq |S_M| \left( 8^p + \sum_{k=1}^m \left( \frac{4}{|k^2 - 2|} \right)^p \right) \leq |S_M| \left( 8^p + \sum_{k=1}^m \frac{4^p}{|k^2 - 2|} \right) \\ &= M(8^p + 8^p) \leq 16^p M. \end{aligned}$$

Observe that each term in  $P_n$  has a zero at 1 with multiplicity at least  $m + 1 > \lfloor \sqrt{n/M} \rfloor$  zeros at 1, and hence so does  $P_n$ .  $\square$

### Proof of Lemma 3.2.

The statement follows from Lemma 3.1 by choosing  $1 \leq M$  so that

$$L := \frac{3}{16\pi^2} M^{1-1/p} = \frac{3}{16\pi^2} M^{1/q}.$$

This can be done when  $\frac{3}{16\pi^2} \leq L$ .  $\square$

### Proof of Lemma 3.3

Let  $L \in (0, 1/17]$ . We define

$$k := \min \left\{ \left\lfloor \frac{-\log L}{\log 17} \right\rfloor, n \right\} \quad \text{and} \quad m := \lfloor \sqrt{n/k} \rfloor.$$

Observe that  $k \geq 1$  and  $m \geq 1$  hold. Let  $P_n := LH_m^k \in \mathcal{P}_n$ , where  $H_m \in \mathcal{P}_{m^2}$  defined by (4.1). Then

$$P_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad j = 0, 1, \dots, n,$$

has at least

$$km \geq k \frac{1}{2} \sqrt{n/k} = \frac{1}{2} \sqrt{nk} = \frac{1}{2\sqrt{\log 17}} \min \left\{ \sqrt{n(-\log L)}, n \right\}$$

zeros at 1, where  $2\sqrt{\log 17} < 7/2$ . Clearly,  $a_{0,n} = P_n(0) = L$ , and using the notation in (4.1), we can deduce that

$$\begin{aligned} \sum_{j=1}^n |a_{j,n}|^p &\leq L^p \left( \sum_{j=1}^n |a_{j,n}| \right)^p \leq L^p \left( 1 + |d_m| + \sum_{k=1}^m |c_{k,m}| \right)^{kp} \\ &\leq L^p (1 + 8 + 8)^{kp} = L^p 17^{kp} \leq L^p L^{-p} = 1, \end{aligned}$$

if  $m \geq 2$ , and

$$\sum_{j=1}^n |a_{j,n}|^p \leq L^p \left( \sum_{j=1}^n |a_{j,n}| \right)^p \leq L^p 2^{kp} \leq L^p L^{-p} = 1,$$

if  $m = 1$ . □

**Proof of Lemma 3.4**

Let

$$r := \left\lfloor 12 \frac{1+L}{1-L} \right\rfloor + 1 \quad \text{and} \quad m := \left\lfloor \sqrt{\frac{n-1}{r}} \right\rfloor.$$

When  $m \leq 1$  we have  $\lfloor (1/9)\sqrt{n(1-L)} \rfloor = 0$ , so there is nothing to prove.

Now assume that  $m \geq 2$ . Let  $P_n \in \mathcal{P}_n$  be defined by  $P_n(x) := H_m(x^r)$ , where  $H_m \in \mathcal{P}_{m^2}$  defined by (4.1). Let  $Q_n \in \mathcal{P}_n$  be defined by

$$Q_n(x) = - \int_0^1 P_n(t) dt + \int_0^x P_n(t) dt.$$

Then, using the notation in (4.1), we have

$$Q_n(x) = -1 - \frac{d_m}{2r+1} - \sum_{k=1}^m \frac{c_{k,m}}{rk^2+1} + x + \frac{d_m x^{2r+1}}{2r+1} + \sum_{k=1}^m \frac{c_{k,m} x^{rk^2+1}}{rk^2+1}.$$

Writing

$$Q_n(x) = \sum_{j=0}^n a_{j,n} x^j, \quad a_{j,n} \in \mathbb{R}, \quad j = 0, 1, \dots, n,$$

and recalling (4.2) and (4.3), we have

$$|a_{0,n}| \geq 1 - \frac{8}{2r+1} - \sum_{k=1}^m \frac{4}{|k^2-4|(rk^2+1)} \geq 1 - \frac{8}{2r+1} - \frac{8}{r} > 1 - \frac{12}{r},$$

and

$$\sum_{j=1}^n |a_{j,n}| \leq 1 + \frac{8}{2r+1} + \sum_{k=1}^m \frac{4}{(k^2-2)(rk^2+1)} < 1 + \frac{12}{r}.$$

Combining the previous two inequalities, we obtain

$$\frac{|a_{0,n}|}{\sum_{j=1}^n |a_{j,n}|} > \frac{1-12/r}{1+12/r} \geq \frac{1-(1-L)/(1+L)}{1+(1-L)/(1+L)} = L.$$

Also  $Q_n$  has at least  $m+1 \geq \lfloor \sqrt{(n-1)/r} \rfloor + 1 \geq \frac{1}{5} \sqrt{(n-1)(1-L)}$  zeros at 1. □

**Proof of Lemma 3.6.**

We assume that  $p, q \in (1, \infty)$ , the result in the cases  $p = 1, q = \infty$  and  $p = \infty, q = 1$  can be proved similarly with straightforward modification of the proof. Let  $m := \mu_q(n, L)$ . Let  $Q$  be a polynomial of degree  $m$  with complex coefficients such that

$$|Q(0)| > \frac{1}{L} \left( \sum_{j=1}^n |Q(j)|^q \right)^{1/q}.$$

Now let  $P$  be a polynomial of the form

$$P(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| \geq L \left( \sum_{j=1}^n |a_j|^p \right)^{1/p}, \quad a_j \in \mathbb{C}.$$

It follows from Hölder's inequality that

$$\left| \sum_{j=1}^n a_j Q(j) \right| \leq \left( \sum_{j=1}^n |a_j|^p \right)^{1/p} \left( \sum_{j=1}^n |Q(j)|^q \right)^{1/q} < \frac{|a_0|}{L} L |Q(0)| = |a_0 Q(0)|.$$

Then  $\sum_{j=0}^n a_j Q(j) \neq 0$ , and hence Lemma 3.5 implies that  $(x-1)^{m+1}$  does not divide  $P$ . We conclude that  $\kappa_p(n, L) \leq m = \mu_q(n, L)$ .  $\square$

**Proof of Lemma 3.9.**

Note that  $\mu_q(n, K) \leq n$  for all  $n \in \mathbb{N}$  and  $L > 0$ , as it is shown by  $H \in \mathcal{P}_n$  defined by  $H(x) := \prod_{j=1}^n (x-j)$ .

Case 1:  $0 < K < n^{-1/q}$ . The choice  $F \equiv 1$  gives the lemma.

Case 2:  $n^{-1/q} \leq K < 2$ . Let  $F$  be the polynomial given in Lemma 3.7 with  $A := n$  and  $M := (4K)^{2q}$ . Then

$$\begin{aligned} \sum_{j=1}^n |F(j)|^q &\leq \sum_{j \leq M^{-1/2}} M^q + \sum_{j > M^{1/2}} \frac{1}{j^{2q}} < M^{q-1/2} + \frac{1}{2q-1} [M^{-1/2}]^{-2q+1} \\ &\leq (1 + 2^{2q-1}) M^{q-1/2}, \end{aligned}$$

so

$$\left( \sum_{j=1}^n |F(j)|^q \right)^{1/q} < 4M^{1-1/(2q)} = K^{-1} F(0),$$

and the degree  $m$  of  $F$  satisfies

$$m < \pi \sqrt{n} \sqrt[4]{M} + 2 < \pi \sqrt{n} (4K)^{q/2} + 2 \leq \sqrt{n} (40K)^{q/2} + 2.$$

Case 3:  $2 \leq K \leq \exp(n - 2\sqrt{n})$ . Let  $R := \lfloor \sqrt{\log K} \rfloor + 1$ , and let  $F$  be the polynomial given in Lemma 3.7 with this  $R$ . Then

$$|F(0)| > K \sum_{j=1}^n |F(j)| \geq K \left( \sum_{j=1}^n |F(j)|^q \right)^{1/q},$$

and the degree  $m$  of  $F$  satisfies

$$m \leq \frac{44}{7} R \sqrt{n} + 4 \leq 13 \sqrt{n \log K} + 4.$$

Case 4:  $K > \exp(n - 2\sqrt{n})$ ,  $n \geq 9$ . Then  $\log K > n - 2\sqrt{n} \geq n/3$  for all  $n \geq 9$ . Hence the polynomial  $F \in \mathcal{P}_n$  defined by  $F(x) := \prod_{j=1}^n (x - j)$  shows that

$$\mu_q(n, K) \leq n \leq \sqrt{3} \min \left\{ \sqrt{n \log K}, n \right\}.$$

Case 5:  $K \geq 2$  and  $n < 9$ . Now the polynomial  $F \in \mathcal{P}_n$  defined by  $F(x) := \prod_{j=1}^n (x - j)$  shows

$$\mu_q(n, K) \leq n \leq 4 \min \left\{ \sqrt{n \log K}, n \right\}.$$

□

### Proof of Lemma 3.10.

First let  $1 < K < 2$ . Let  $m = \lfloor \sqrt{n(K-1)/2} \rfloor + 1$ . Let  $T_m$  be the Chebyshev polynomial of degree  $m$  defined by

$$T_m(\cos t) = \cos(mt), \quad t \in \mathbb{R}.$$

It is well known that  $|T'_m(1)| = m^2$  and  $T'_m(x)$  is increasing on  $[1, \infty)$ , hence  $T_m(1+x) \geq 1 + m^2x$  for all  $x > 0$ . Now we define  $F \in \mathcal{P}_m$  by

$$F(x) := T_m \left( \frac{-2x}{n-1} + \frac{n+1}{n-1} \right).$$

Then  $|F(x)| \leq 1$  for all  $x \in [1, n]$ , and

$$F(0) \geq T_m \left( 1 + \frac{2}{n-1} \right) > 1 + \frac{m^2}{n-1} > 1 + \frac{m^2}{n} \geq K,$$

which finishes the proof in the case of  $1 < K < 2$ . Now let  $k \geq 2$ . Then the polynomial  $F \in \mathcal{P}_m$  chosen for  $q = 1$ ,  $n \in \mathbb{N}$ , and  $K \geq 2$  by Lemma 3.9 gives that

$$|F(0)| > K \left( \sum_{j=1}^n |F(j)|^q \right)^{1/q} \geq K \max_{j \in \{1, 2, \dots, n\}} |F(j)|,$$

with

$$m \leq 13 \min \left\{ \sqrt{n \log K}, n \right\} + 4.$$

□

## 5 Proofs of the Theorems.

### Proof of Theorem 2.1.

Without loss of generality we may assume that  $p \in (1, \infty)$ , as the case

$p = \infty$  follows by a simple limiting argument (or we may as well refer to the main result in [B-13]). By Lemma 3.6 we have

$$\kappa_p(n, L) \leq \mu_q(n, L)$$

for every  $n \in \mathbb{N}$  and  $L > 0$ . The lower bounds for  $\kappa_p(n, L)$  follows from Lemmas 3.2 and 3.3. The upper bounds for  $\mu_q(n, L)$  follow from Lemma 3.9 with  $K = L^{-1}$ .  $\square$

### Proof of Theorem 2.2.

By Lemma 3.6 we have

$$\kappa_1(n, L) \leq \mu_\infty(n, L)$$

for every  $n \in \mathbb{N}$  and  $L > 0$ . The lower bounds for  $\kappa_1(n, L)$  follow from Lemmas 3.3 and 3.4. The upper bounds for  $\mu_\infty(n, L)$  follow from Lemma 3.10 with  $K = L^{-1}$ .  $\square$

## 6 Remarks and Problems.

A question that we have not really considered in this paper is the following. Are here are examples of  $n$ ,  $L$ , and  $p$  for which the values of  $\kappa_p(n, L)$  are significantly smaller if the coefficients are required to be rational (perhaps together with other restrictions). The same question may be raised about  $\mu_q(n, L)$ . As the conditions on the coefficients of the polynomials in Theorems 2.1 and 2.2 are homogeneous, assuming rational coefficients and integer coefficients lead to the same results. Three special classes of interest are

$$\mathcal{F}_n := \left\{ Q : Q(z) = \sum_{j=0}^n a_j z^j, \ a_j \in \{-1, 0, 1\} \right\},$$

$$\mathcal{L}_n := \left\{ Q : Q(z) = \sum_{j=0}^n a_j z^j, \ a_j \in \{-1, 1\} \right\},$$

and

$$\mathcal{K}_n := \left\{ Q : Q(z) = \sum_{j=0}^n a_j z^j, \ a_j \in \mathbb{C}, \ |a_j| = 1 \right\}.$$

The following three problems arise naturally.

**Problem 6.1.** *How many zeros can a polynomial  $0 \neq Q \in \mathcal{F}_n$  have at 1?*

**Problem 6.2.** *How many zeros can a polynomial  $Q \in \mathcal{L}_n$  have at 1?*

**Problem 6.3.** *How many zeros can a polynomial  $Q \in \mathcal{K}_n$  have at 1?*

The case  $p = \infty$  and  $L = 1$  in our Theorem 2.1 gives that every  $0 \neq Q \in \mathcal{F}_n$ , every  $Q \in \mathcal{L}_n$ , and every  $Q \in \mathcal{K}_n$  can have at most  $cn^{1/2}$  zeros at 1, where  $c > 0$  is an absolute constant. However, one may expect better results by utilizing the additional pieces of information on their coefficients.

It was observed in [B-99] that for every integer  $n \geq 2$  there is a  $Q \in \mathcal{F}_n$  having at least  $c(n/\log n)^{1/2}$  zeros at 1 with an absolute constant  $c > 0$ . This can be shown by a simple pigeon hole argument. However, as far as we know, closing the gap between  $cn^{1/2}$  and  $c(n/\log n)^{1/2}$  in Problem 6.1 is an open and most likely a very difficult problem.

As far as Problem 6.2 is concerned, Boyd [B-97] showed that for  $n \geq 3$  every  $Q \in \mathcal{L}_n$  has at most

$$(6.1) \quad \frac{c(\log n)^2}{\log \log n}$$

zeros at 1. This is the best known upper bound in Problem 6.2 even today. Boyd's proof is very clever and, up to an application of the Prime Number Theorem, completely elementary. It is reasonable to conjecture that there is an absolute constant  $c > 0$  such that every  $Q \in \mathcal{L}_n$ ,  $n \geq 2$ , has at most  $c \log n$  zeros at 1 for every  $n$ . It is easy to see that for every integer  $n \geq 2$  there are  $Q_n \in \mathcal{L}_n$  with at least  $c \log n$  zeros at 1 with an absolute constant  $c > 0$ .

As far as Problem 6.3 is concerned, one may suspect that every every  $Q \in \mathcal{K}_n$ ,  $n \geq 2$ , has at most  $c \log n$  zeros at 1. However, just to see if Boyd's bound (6.1) holds for every  $Q \in \mathcal{K}_n$  seems quite challenging and beyond reach at the moment.

## References

- [A-90] F. Amoroso *Sur le diamètre transfini entier d'un intervalle réel*, Ann. Inst. Fourier, Grenoble 40 (1990), 885–911.
- [A-02] V.V. Andrievskii and H-P. Blatt *Discrepancy of Signed Measures and Polynomial Approximation*, Springer, New York, 2002.
- [A-79] E. Aparicio Bernardo *New bounds on the minimal Diophantine deviation from zero on  $[0, 1]$  and  $[0, 1/4]$* , Proceedings of the sixth conference of Portuguese and Spanish mathematicians, Part I, Santander (1979), 289–291.

- [B-98] F. Beaucoup, P. Borwein, D.W. Boyd and C. Pinner *Multiple roots of  $[-1, 1]$  power series*, J. London Math. Soc. 2 57 (1998), 135–147.
- [B-32] A. Bloch and G. Pólya *On the roots of certain algebraic equations*, Proc. London Math. Soc 33 (1932), 102–114.
- [B-87] E. Bombieri and J. Vaaler *Polynomials with low height and prescribed vanishing*, Analytic Number Theory and Diophantine Problems, Birkhäuser 1987, 53–73.
- [B-02] P. Borwein *Computational Excursions in Analysis and Number Theory*, Springer, New York, 2002.
- [B-95] P. Borwein and T. Erdélyi *Polynomials and Polynomial Inequalities*, Springer, New York, 1995.
- [B-96] P. Borwein and T. Erdélyi *The integer Chebyshev problem*, Math. Comp. 65 (1996), 661–681.
- [B-97a] P. Borwein and T. Erdélyi *On the zeros of polynomials with restricted coefficients*, Illinois J. Math. 41 (1997), 667–675.
- [B-97b] P. Borwein and T. Erdélyi *Generalizations of Müntz’s theorem via a Remez-type inequality for Müntz spaces*, J. Amer. Math. Soc. 10 (1997), 327–349.
- [B-07] P. Borwein and T. Erdélyi *Lower bounds for the number of zeros of cosine polynomials: a problem of Littlewood*, Acta Arith. 128 (2007), 377–384.
- [B-08a] P. Borwein, T. Erdélyi, R. Ferguson, and R. Lockhart *On the zeros of cosine polynomials : solution to a problem of Littlewood*, Ann. of Math. (2) 167 (2008), 1109–1117.
- [B-99] P. Borwein, T. Erdélyi and G. Kós *Littlewood-type problems on  $[0, 1]$* , Proc. London Math. Soc. [79] (1999), 22–46.
- [B-13] P. Borwein, T. Erdélyi and G. Kós *The multiplicity of the zero at 1 of polynomials with constrained coefficients*, Acta Arith. 159 (2013) no. 4, 387–395.
- [B-08b] P. Borwein, T. Erdélyi and F. Littmann *Polynomials with coefficients from a finite set*, Trans. Amer. Math. Soc. 360 (2008), 5145–5154.

- [B-92] P. Borwein, T. Erdélyi and J. Zhang *Müntz systems and orthogonal Müntz-Legendre polynomials*, Trans. Amer. Math. Soc. 342 (1992), 523–542.
- [B-94] P. Borwein and C. Ingalls *The Prouhet-Tarry-Escott problem revisited*, Ens. Math. 40 (1994), 3–27.
- [B-00] P. Borwein and M.J. Mossinghoff *Polynomials with height 1 and prescribed vanishing at 1*, Experiment. Math. 9 (2000), 425–433.
- [B-97] D.W. Boyd *On a problem of Byrnes concerning polynomials with restricted coefficients*, Math. Comp. 66 (1997), 1697–1703.
- [Bu-99] H. Buhrman, R. Cleve, R. de Wolf, and C. Zalka *Bounds for small-error and zero-error quantum algorithms* 40th Annual Symposium on Foundations of Computer Science, New York, 1999, 358–368, IEEE Computer Soc., Los Alamitos, CA.
- [C-02] P.G. Casazza and N.J. Kalton *Roots of complex polynomials and Weyl-Heisenberg frame sets*, Proc. Amer. Math. Soc. 130 (2002) no. 8, 2313–2318.
- [C-13] J.M. Cooper and A.M. Dutele *Greedy Galois games*, Amer. Math. Monthly 120 (2013) no. 5, 441451.
- [C-92] D. Coppersmith and T.J. Rivlin *The growth of polynomials bounded at equally spaced points* SIAM J. Math. Anal. 23 (1992) no. 4, 970–983.
- [C-10] E. Croot and D. Hart  *$h$ -fold sums from a set with few products*, SIAM J. Discrete Math. 24 (2010) no. 2, 505–519.
- [D-99] A. Dubickas *On the order of vanishing at 1 of a polynomial*, Lithuanian Math. J. 39 (1999), 365–370.
- [D-01] A. Dubickas *Three problems of polynomials of small measure*, Acta Arith. 98 (2001), 279–292.
- [D-13] A. Dubickas *Polynomials with multiple roots at 1*, Int. J. Number Theory 10 (2014) no. 2, 391–400.
- [D-03] M. Dudik and L.J. Schulman *Reconstruction from subsequences*, Journal of Combinatorial Theory, Series A 103 (2003) no. 2, 337–348.



- [E-02a] T. Erdélyi *Markov-Bernstein type inequalities for polynomials under Erdős-type constraints*, Paul Erdős and his Mathematics I, Bolyai Society Mathematical Studies, 11, Gábor Halász, László Lovász, Dezső Miklós, and Vera T. Sós (Eds.), Springer Verlag, New York, 2002, 219–239.
- [E-08a] T. Erdélyi Extensions of the Bloch-Pólya theorem on the number of real zeros of polynomials, *Journal de théorie des nombres de Bordeaux* 20 (2008), 281–287.
- [E-02b] T. Erdélyi *Polynomials with Littlewood-type coefficient constraints*, Approximation Theory X: Abstract and Classical Analysis, Charles K. Chui, Larry L. Schumaker, and Joachim Stöckler (Eds.), Vanderbilt University Press, Nashville, TN, 2002, 153–196.
- [E-08b] T. Erdélyi *An improvement of the Erdős-Turán theorem on the zero distribution of the zeros of polynomials*, *C. R. Acad. Sci. Paris Sér. I Math.* 346 (2008), 267–270.
- [E-13] T. Erdélyi *Pseudo-Boolean functions and the multiplicity of the zeros of polynomials*, *Journal d'Analyse Math.* 127, (2015), 91–108.
- [E-50] P. Erdős and P. Turán *On the distribution of roots of polynomials*, *Annals of Math.* 57 (1950), 105–119.
- [F-80] Le Baron O. Ferguson *Approximation by Polynomials with Integral Coefficients*, Amer. Math. Soc., Rhode Island, 1980.
- [F-00] W. Foster and I. Krasikov *An improvement of a Borwein-Erdélyi-Kós result*, *Methods Appl. Anal.* 7 (2000) no. 4, 605–614.
- [G-05] C.S. Güntürk *Approximation by power series with  $\pm 1$  coefficients*, *Int. Math. Res. Not.* (2005) no. 26, 1601–1610.
- [H-38] G.H. Hardy and E.M. Wright *An Introduction to the Theory of Numbers*, Clarendon Press, Oxford, 1938.
- [H-82] L.K. Hua *Introduction to Number Theory*, Springer-Verlag, Berlin Heidelberg, New York, 1982.
- [K-03] G. Kós, P. Ligeti, and P. Sziklai *Reconstruction of matrices from submatrices*, *Math. Comp.* 78 (2009), 1733–1747.

- [K-04] I. Krasikov *Multiplicity of zeros and discrete orthogonal polynomials*, Results Math. 45 (2004), 1–2, 59–66.
- [M-68] M. Minsky and S. Papert *Perceptrons: An Introduction to Computational Geometry*, MIT Press, Cambridge, Mass., 1968.
- [M-03] M.J. Mossinghoff *Polynomials with restricted coefficients and prescribed noncyclotomic factors*, London Math. Soc. J. Comput. Math. 6 (2003), 314–325 (electronic).
- [N-94] N. Nisan and M. Szegedy *On the degree of Boolean functions as real polynomials*, Computational Complexity 4 (1994) no. 4, 301–313, Earlier version in STOC92.
- [O-93] A.M. Odlyzko and B. Poonen *Zeros of polynomials with 0, 1 coefficients*, Enseign. Math. (2) 39 (1993), 317–348.
- [P-99] C. Pinner *Double roots of  $[-1, 1]$  power series and related matters*, Math. Comp. 68 (1999), 1149–1178.
- [P-13] I.E. Pritsker and A.A. Sola *Expected discrepancy for zeros of random algebraic polynomials*, Proc. Amer. Math. Soc. 142 (2014), 4251–4263.
- [R-07] E.A. Rakhmanov *Bounds for polynomials with a unit discrete norm* Ann. of Math. 165 (2007), 55–88.
- [R-04] F. Rodier *Sur la non-linéarité des fonctions booléennes* Acta Arith. 115 (2004) no. 1, 1–22.
- [Sch-33] I. Schur *Untersuchungen über algebraische Gleichungen*, Sitz. Preuss. Akad. Wiss., Phys.-Math. Kl. (1933), 403–428.
- [S-99] I.E. Shparlinski *Finite Fields: Theory and Computation: the Meeting Point of Number Theory, Computer Science, Coding Theory and Cryptography*, Dordrecht, London, 1999.
- [Sz-34] G. Szegő *Bemerkungen zu einem Satz von E. Schmidt über algebraische Gleichungen* Sitz. Preuss. Akad. Wiss., Phys.-Math. Kl. (1934), 86–98.
- [T-07] V. Totik and P. Varjú *Polynomials with prescribed zeros and small norm*, Acta Sci. Math. (Szeged) 73 (2007) no. 3–4, 593–611.
- [T-84] P. Turán *On a New Method of Analysis and its Applications*, Wiley, New York, 1984.