

MARKOV AND BERNSTEIN TYPE INEQUALITIES ON SUBSETS OF $[-1, 1]$ AND $[-\pi, \pi]$

PETER BORWEIN AND TAMÁS ERDÉLYI

ABSTRACT. We extend Markov's, Bernsteins's, and Videnskii's inequalities to arbitrary subsets of $[-1, 1]$ and $[-\pi, \pi]$, respectively.

The primary purpose of this note is to extend Markov's and Bernsteins's inequalities to arbitrary subsets of $[-1, 1]$ and $[-\pi, \pi]$, respectively.

We denote by \mathcal{P}_n the set of all real algebraic polynomials of degree at most n and let $m(\cdot)$ denote the Lebesgue measure of a subset of \mathbb{R} . We were led to the results of this paper by the following problem. Can one give polynomials $p_n \in \mathcal{P}_n$ and numbers $a_n \in (0, 1)$, $n = 1, 2, \dots$, such that

$$(i) \quad m(\{x \in [0, 1] : |p_n(x)| \leq 1\}) \geq 1 - a_n,$$

$$(ii) \quad \max_{0 \leq x \leq a_n} |p_n(x)| \leq 1$$

and

$$(iii) \quad \lim_{n \rightarrow \infty} n^{-2} |p'_n(0)| = \infty$$

are satisfied? This question was asked by Vilmos Totik, and a positive answer would have been used in proving a conjecture in the theory of orthogonal polynomials. However, Theorem 2 of this note shows that the answer to the above question is negative, in fact, it gives slightly more. In addition, our Theorem 1 answers the corresponding question for trigonometric polynomials. Though our results cannot be used for Totik's original purpose, our proofs illustrate well, how Remez-type inequalities can be used in proving various other polynomial inequalities.

In this note we prove the following pair of theorems.

Theorem 1. *Let $0 < a \leq 2\pi$, $0 < L \leq 1$, let A be a closed subset of $[0, 2\pi]$ with Lebesgue measure $m(A) \geq 2\pi - a$. There is an absolute constant $c_1 > 0$ such that*

$$(1) \quad \max_{t \in I} |p'(t)| \leq c_1 L^{-1} (n + n^2 a) \max_{t \in A} |p(t)|$$

for every real trigonometric polynomial p of degree at most n and for every subinterval I of A with length at least La .

This material is based upon work supported by the National Science and Engineering Research Council of Canada (P.B.) and the National Science Foundation under Grant No. DMS-9024901 (T.E.)

Theorem 2. *Let $0 < a \leq 1$, $0 < M \leq 1$, let A be a closed subset of $[0, 1]$ with Lebesgue measure $m(A) \geq 1 - a$. There is an absolute constant $c_2 > 0$ such that*

$$(2) \quad \max_{x \in I} |p'(x)| \leq c_2 M^{-1} n^2 \max_{x \in A} |p(x)|$$

for every real algebraic polynomial p of degree at most n and for every subinterval I of A with length at least Ma .

Up to the constant c_1 , Theorem 1 is an extension of both Bernstein's [5, pp. 39-41] and Videnskii's [6] inequalities, while up to the constant c_2 , Theorem 2 contains Markov's inequality [5, pp. 39-41] as a special case.

The key to the proof of Theorem 1 is a Remez-type inequality [2] proved recently for trigonometric polynomials, while the proof of Theorem 2 relies on Theorem 1.

Proof of Theorem 1. Denote by \mathcal{T}_n the set of all trigonometric polynomials of degree at most n with real coefficients. If $\pi/2 \leq a \leq 2\pi$, then the theorem follows from an extension [1, Theorem 5] of an inequality of Videnskii [6]. Therefore, in the sequel we assume that $0 < a < \pi/2$. Let I be a subinterval of A such that $m(I) \geq La$ and $\pi \in I$. It is sufficient to prove that there is an absolute constant $c_1 > 0$ such that

$$(3) \quad |p'(\pi)| \leq c_1 L^{-1} (n + n^2 a) \max_{t \in A} |p(t)|$$

for every $p \in \mathcal{T}_n$. Let T_n be the Chebyshev polynomial of degree n given by

$$(4) \quad T_n(x) = \cos(n \arccos x), \quad -1 \leq x \leq 1,$$

and let

$$(5) \quad Q_{n,La}(t) := T_{2n}(\sin(t/2)(\cos(La/4))^{-1})(T_{2n}((\cos(La/4))^{-1}))^{-1}.$$

A simple calculation shows that

$$(6) \quad Q_{n,La}(\pi) = 1, \quad Q'_{n,La}(\pi) = 0, \quad \max_{t \in \mathbb{R}} |Q_{n,La}(t)| = 1,$$

and there is an absolute constant $c_3 > 0$ such that

$$(7) \quad |Q_{n,La}(t)| \leq \exp(-c_3 n La), \quad t \in [0, \pi - La/2] \cup [\pi + La/2, 2\pi].$$

Let $p \in \mathcal{T}_n$ be such that

$$(8) \quad \max_{t \in A} |p(t)| = 1.$$

The Remez-type inequality for trigonometric polynomials [2, Theorem 2], $m(A) \geq 2\pi - a$, $0 < a \leq \pi/2$, and (8) yield that there is an absolute constant $c_4 > 0$ such that

$$(9) \quad \max |p(t)| < \exp(c_4 n a).$$

Denote the endpoints of the interval I by $\alpha < \beta$. Since $\beta - \alpha = m(I) \geq La$ and $\pi \in I$, we have either $\alpha \leq \pi - La/2$ or $\beta \geq \pi + La/2$. We may assume that

$$(10) \quad \beta \geq \pi + La/2,$$

otherwise we consider the trigonometric polynomial $\tilde{p} \in \mathcal{T}_n$ defined by $\tilde{p}(t) := p(\pi - t)$. Now let

$$(11) \quad m := [c_4 c_3^{-1} L^{-1} n] + 1 \quad \text{and} \quad Q := Q_{m, La}.$$

Observe that (6) - (11) imply

$$(12) \quad |(pQ)(t)| \leq 1, \quad t \in E,$$

where

$$(13) \quad E := [0, \pi - La/2] \cup [\pi, 2\pi].$$

Note that E is an interval of the period with length $2\pi - La/2$, and $\pi \in E$. Therefore an extension [1, Theorem 5] of an inequality of Videnskii [6], $0 < L \leq 1$ and (8) yield that there are absolute constants $c_5 > 0$ and $c_1 > 0$ such that

$$(14) \quad \begin{aligned} |(pQ)'(\pi)| &\leq ((n+m) + c_5(n+m)^2 La/2) \\ &\leq c_1 L^{-1} (n + n^2 a) \max_{t \in A} |p(t)|. \end{aligned}$$

Recalling (6), we have

$$(15) \quad p'(\pi) = (pQ)'(\pi),$$

which, together with (14) gives the theorem. ■

Proof of Theorem 2. If $1/4 \leq a \leq 1$, then the theorem follows from the Markov inequality [5, pp. 39-41]. Therefore, in what follows we may assume that $0 < a \leq 1/4$. Without loss of generality we may assume that $I = [0, b]$, where $Ma \leq b \leq 1$, the general case can be deduced from this easily by a linear transformation. Let $p \in \mathcal{P}_n$,

$$(16) \quad y(t) := 1/2 + (1/2 + a) \cos t,$$

$$(17) \quad \tilde{p}(t) := p(y(t)) \in \mathcal{T}_n$$

$$(18) \quad \tilde{A} := \{t \in [0, 2\pi] : y(t) \in A\},$$

$$(19) \quad \tilde{I} := \{t \in [0, \pi] : y(t) \in I\}$$

and

$$(20) \quad \tilde{a} := 2\pi - m(\tilde{A}), \text{ i.e. } m(\tilde{A}) = 2\pi - \tilde{a}.$$

It is easy to see that $0 < a \leq 1/4$, $A \subset [0, 1]$, $m(A) \geq 1 - a$, $m(I) \geq Ma$, (16), (18), (19), and (20) imply that

$$(21) \quad \tilde{a} \leq \dots \sqrt{\dots}$$

and

$$(22) \quad m(\tilde{I}) \geq c_7 M \sqrt{a} \geq c_7 c_6^{-1} M \tilde{a}$$

with suitable absolute constants $c_6 > 0$ and $c_7 > 0$. If $L := c_7 c_6^{-1} M \leq 1$ and $a \geq n^{-2}$, then Theorem 1, (20), (21) and (22) yield

$$(23) \quad \begin{aligned} \max_{t \in \tilde{I}} |\tilde{p}'(t)| &\leq c_1 c_7^{-1} c_6 M^{-1} (n + n^2 \tilde{a}) \max_{t \in \tilde{A}} |\tilde{p}(t)| \\ &\leq c_8 M^{-1} n^2 \sqrt{a} \max_{x \in A} |p(x)| \end{aligned}$$

with a suitable absolute constant $c_8 > 0$. Also, (16) - (19) and $I \subset [0, 1]$ imply that

$$|\tilde{p}'(t)| = |p'(y(t))y'(t)| = |p'(y(t))|(1/2 + a) \sin t \geq c_9 |p'(y(t))| \sqrt{a}$$

for every $t \in \tilde{I}$ with a suitable absolute constant $c_9 > 0$. Since every $x \in I$ is of the form $x = y(t)$ with some $t \in \tilde{I}$, (23) and (24) imply that

$$(25) \quad \max_{x \in I} |p'(x)| \leq c_8 c_9^{-1} M^{-1} n^2 \max_{x \in A} |p(x)|,$$

whenever $c_7 c_6^{-1} M \leq 1$ and $a \geq n^{-2}$. If $c_7 c_6^{-1} M \geq 1$, i.e. $M \geq c_6 c_7^{-1}$, and $a \geq n^{-2}$, then I can be divided into subintervals of length $k^{-1}m(I)$, where $k := \lceil c_6 c_7^{-1} \rceil + 1$, and the already proved part gives the theorem. If $0 < a < n^{-2}$, $A \subset [0, 1]$ and $m(A) \geq 1 - a$, then the Remez inequality [4, p. 119-121] or [3] yields that

$$(26) \quad \max_{0 \leq x \leq 1} |p(x)| \leq c_{10} \max_{x \in A} |p(x)|$$

for every $p \in \mathcal{P}_n$, where $c_{10} > 0$ is a suitable absolute constant. Combining this with the Markov inequality [5, p. 39-41], we obtain

$$(27) \quad \begin{aligned} \max_{x \in I} |p'(x)| &\leq \max_{0 \leq x \leq 1} |p'(x)| \leq 2n^2 \max_{0 \leq x \leq 1} |p(x)| \\ &\leq 2c_{10} n^2 \max_{x \in A} |p(x)|, \end{aligned}$$

and the theorem is completely proved. ■

It may be interesting to compare Theorem 2 with the following

Example 3. Let $0 < a \leq 1/2$, $A = [0, 1 - a] \cup \{1\}$ and

$$P_n(x) = (x - 1)T_n(2(1 - a)^{-1}x - 1), \quad n = 1, 2, \dots,$$

where T_n is the Chebyshev polynomial of degree n defined by $T_n(x) = \cos(n \arccos x)$, $-1 \leq x \leq 1$. Then

$$\begin{aligned} \max_{x \in A} |P_n'(x)| &\geq |P_n'(1)| = T_n(2(1 - a)^{-1} - 1) \geq T_n(1 + 2a) \\ &\geq 2^{-1}(1 + 2\sqrt{a})^n \geq 2^{-1}(1 + 2\sqrt{a})^n \max_{x \in A} |P_n(x)|. \end{aligned}$$

A similar example can be given in the trigonometric case.

REFERENCES

- [1] T. Erdélyi, *Markov-type estimates for derivatives of polynomials of special type*, Acta. Math. Hung. **51** (1988), 421-436.
- [2] ———, *Remez-type inequalities on the size of generalized polynomials*, J. London Math. Soc. (to appear).
- [3] ———, *The Remez inequality on the size of polynomials*, “Approximation Theory VI, Vol I”, C. K. Chui, L. L. Schumaker and J. D. Ward, Eds., Academic Press, Boston, 1989, 243-246.
- [4] G. Freud, “*Orthogonal Polynomials*”, Pergamon Press, Oxford, 1971.
- [5] G. G. Lorentz, “*Approximation of Functions*”, Holt, Rinehart and Winston, New York 1966.
- [6] V. S. Videnskii, *Extremal estimates for the derivative of a trigonometric polynomial on an interval shorter than the period*, (in Russian), Dokl. Akad. Nauk. USSR **130** (1960), 13-16.

Acknowledgement. The authors wish to thank Vilmos Totik for his question that helped to formulate the theorems of this paper.

DEPARTMENT OF MATHEMATICS DALHOUSIE UNIVERSITY, HALIFAX, NOVA SCOTIA,
CANADA B3H 3J5

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OHIO 43210,
USA