

Name _____

MATH 251 Exam 1 Version B Fall 2017
Sections 515 Solutions P. Yasskin

1-9	/54	11	/16
10	/33	Total	/103

Multiple Choice: (6 points each. No part credit.)

1. The points $A = (2, -3, 4)$ and $B = (4, 1, 0)$ are the endpoints of the diameter of a sphere. What is the radius of the sphere?

- a. 2
- b. 3 Correct Choice
- c. 4
- d. 5
- e. 6

Solution: The diameter is $d = d(A, B) = \sqrt{(4-2)^2 + (1-(-3))^2 + (0-4)^2} = \sqrt{4+16+16} = 6$.
The radius is $r = 3$.

2. The points $A = (2, -3, 4)$ and $B = (4, 1, 0)$ are the endpoints of the diameter of a sphere. What is the center of the sphere?

- a. (2, 4, -4)
- b. (3, 1, 2)
- c. (6, 2, 4)
- d. (3, -1, 2) Correct Choice
- e. (6, -2, 4)

Solution: The center is the midpoint of the diameter: $C = \frac{A+B}{2} = (3, -1, 2)$.

3. Find the angle between the normals to the planes $3x + 2y - 4z = 3$ and $2x - y + z = 2$.

- a. 90° Correct Choice
- b. 60°
- c. 45°
- d. 30°
- e. 0°


Solution: The normals are $\vec{N}_1 = (3, 2, -4)$ and $\vec{N}_2 = (2, -1, 1)$.
Since $\vec{N}_1 \cdot \vec{N}_2 = 6 - 2 - 4 = 0$, the vectors are perpendicular.

4. Duke Skywalker pushes an asteroid from the point $P = (2, -3, 5)$ to the point $Q = (5, -1, 4)$ by the force $\vec{F} = (4, 1, 2)$. Find the work done to move the asteroid.
- 2
 - 4
 - 6
 - 12 Correct Choice
 - 16

Solution: The displacement is $\vec{PQ} = Q - P = (3, 2, -1)$. So the work done is $W = \vec{F} \cdot \vec{PQ} = 12 + 2 - 2 = 12$

5. The plot at the right is which polar equation?

- $r = 1 + \cos 3\theta$
- $r = 1 - 2 \cos 3\theta$
- $r = 1 + 2 \cos 3\theta$
- $r = 2 - \cos 3\theta$
- $r = 2 + \cos 3\theta$ Correct Choice



Solution: The rectangular plot is

Further, $r = 3$ when $\theta = 0$ and r never goes negative.

6. Find a vector perpendicular to the plane containing the points $P = (2, 1, 4)$, $Q = (-1, 3, 2)$ and $R = (3, 1, 2)$
- $(2, -1, 2)$
 - $(-4, 8, -2)$
 - $(2, 4, 1)$ Correct Choice
 - $(2, -2, 1)$
 - $(-4, 2, -4)$

Solution: $\vec{PQ} = Q - P = (-3, 2, -2)$ $\vec{PR} = R - P = (1, 0, -2)$

$$\vec{N} = \vec{PQ} \times \vec{PR} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -3 & 2 & -2 \\ 1 & 0 & -2 \end{vmatrix} = \hat{i}(-4 - 0) - \hat{j}(6 + 2) + \hat{k}(0 - 2) = (-4, -8, -2) \text{ or any multiple.}$$

7. If $|\vec{u}| = 2$ and $|\vec{v}| = 5$ and $\vec{u} \cdot \vec{v} = 6$ find $|\vec{u} \times \vec{v}|$.


- a. 0
- b. 2
- c. 4
- d. 6
- e. 8 Correct Choice

Solution: By the Pythagorean Identity for Dot and Cross Products, we have

$$|\vec{u} \times \vec{v}| = \sqrt{|\vec{u}|^2 |\vec{v}|^2 - (\vec{u} \cdot \vec{v})^2} = \sqrt{100 - 36} = 8$$

8. The plot at the right is the graph of which equation?

- a. $x^2 + y^2 - z^2 = -1$ Correct Choice
- b. $x^2 + y^2 - z^2 = 0$
- c. $x^2 + y^2 - z^2 = 1$
- d. $x^2 + y^2 - z = 1$
- e. $x^2 + y^2 - z = -1$



Solution: (e) is $x^2 + y^2 + 1 = z^2$ So $z \geq 1$ or $z \leq -1$.

9. Find the point where the line $(x,y,z) = \vec{r}(t) = (2t + 1, t - 1, 2t - 1)$ intersects the plane $3x + 2y + z = 20$.

At this point $x + y + z =$

- a. 13
- b. 9 Correct Choice
- c. 4
- d. -1
- e. -6

Solution: Plug the line into the plane and solve for t :

$$3x + 2y + z = 3(2t + 1) + 2(t - 1) + (2t - 1) = 10t = 20 \quad \Rightarrow \quad t = 2$$

So the point is $(x,y,z) = \vec{r}(2) = (5, 1, 3)$ and so $x + y + z = 9$.

Work Out: (Points indicated. Part credit possible. Show all work.)

10. (33 points) For the parametric curve $\vec{r}(t) = \left(\frac{2}{t}, 6t, 3t^3\right)$ compute each of the following:

a. (3 pts) velocity \vec{v}

Solution:

$$\vec{v} = \underline{\underline{\left(\frac{-2}{t^2}, 6, 9t^2\right)}}$$

b. (3 pts) acceleration \vec{a}

Solution:

$$\vec{a} = \underline{\underline{\left(\frac{4}{t^3}, 0, 18t\right)}}$$

c. (3 pts) jerk \vec{j}

Solution:

$$\vec{j} = \underline{\underline{\left(\frac{-12}{t^4}, 0, 18\right)}}$$

d. (3 pts) speed $|\vec{v}|$ (Simplify!)

HINT: The quantity inside the square root is a perfect square.

Solution: $|\vec{v}| = \sqrt{\frac{4}{t^4} + 36 + 81t^4} = \sqrt{\left(\frac{2}{t^2} + 9t^2\right)^2}$

$$|\vec{v}| = \underline{\underline{\frac{2}{t^2} + 9t^2}}$$

e. (3 pts) tangential acceleration a_T

Solution: $a_T = \frac{d|\vec{v}|}{dt} = \frac{d}{dt}\left(\frac{2}{t^2} + 9t^2\right)$

$$a_T = \underline{\underline{\frac{-4}{t^3} + 18t}}$$

f. (4 pts) unit binormal \hat{B} (Do this last.)

Solution: $\vec{v} \times \vec{a} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{-2}{t^2} & 6 & 9t^2 \\ \frac{4}{t^3} & 0 & 18t \end{vmatrix} = \hat{i}(108t) - \hat{j}\left(\frac{-36}{t} - \frac{36}{t}\right) + \hat{k}\left(\frac{-24}{t^3}\right)$

$$= \left(108t, \frac{72}{t}, \frac{-24}{t^3}\right) = 12\left(9t, \frac{6}{t}, \frac{-2}{t^3}\right)$$

$$|\vec{v} \times \vec{a}| = 12\sqrt{81t^2 + \frac{36}{t^2} + \frac{4}{t^6}} = 12\left(9t + \frac{2}{t^3}\right) = \frac{12(9t^4 + 2)}{t^3}$$

$$\hat{B} = \frac{\vec{v} \times \vec{a}}{|\vec{v} \times \vec{a}|} = \frac{t^3}{9t^4 + 2}\left(9t, \frac{6}{t}, \frac{-2}{t^3}\right)$$

$$\hat{B} = \underline{\underline{\left(\frac{9t^4}{9t^4 + 2}, \frac{6t^2}{9t^4 + 2}, \frac{-2}{9t^4 + 2}\right)}}$$

Recall: $\vec{r}(t) = \left(\frac{2}{t}, 6t, 3t^3\right)$

g. (2 pts) the values of t where the curve passes thru the points

$A = (2, 6, 3)$

$t = \underline{\quad 1 \quad}$

$B = (1, 12, 24)$

$t = \underline{\quad 2 \quad}$

Solution: Compare each point to the curve $\left(\frac{2}{t}, 6t, 3t^3\right)$. The x component is sufficient, but you should check the other components.

h. (4 pts) arc length between $(2, 6, 3)$ and $(1, 12, 24)$, $L = \int_{(2,6,3)}^{(1,12,24)} ds$

Solution: $L = \int_1^2 |\vec{v}| dt = \int_1^2 \left(\frac{2}{t^2} + 9t^2\right) dt = \left[\frac{-2}{t} + 3t^3\right]_1^2 = (-1 + 24) - (-2 + 3)$

$L = \underline{\quad 22 \quad}$

i. (4 pts) A wire has the shape of this curve between $(2, 6, 3)$ and $(1, 12, 24)$. Find the mass of the wire if the linear mass density is $\rho = \frac{1}{6}xz$.

(Don't simplify the answer.)

Solution: $\vec{v} = \left(\frac{-2}{t^2}, 6, 9t^2\right)$ $|\vec{v}| = \frac{2}{t^2} + 9t^2$ $\rho = \frac{1}{6}xz = \frac{1}{6}\left(\frac{2}{t}\right)(3t^3) = t^2$

$M = \int_{(2,6,3)}^{(1,12,24)} \rho ds = \int_1^2 \frac{1}{6}xz|\vec{v}| dt = \int_1^2 t^2\left(\frac{2}{t^2} + 9t^2\right) dt = \int_1^2 (2 + 9t^4) dt = \left[2t + \frac{9t^5}{5}\right]_1^2$

$M = \underline{\quad \left(4 + \frac{9 \cdot 2^5}{5}\right) - \left(2 + \frac{9}{5}\right) \quad} = \frac{289}{5}$

j. (4 pts) A wire has the shape of this curve. Find the work done by the force $\vec{F} = (z, y, x)$ which pushes a bead along the wire from $(2, 6, 3)$ to $(1, 12, 24)$.

Solution: $\vec{F} = (z, y, x) = \left(3t^3, 6t, \frac{2}{t}\right)$ $\vec{v} = \left(\frac{-2}{t^2}, 6, 9t^2\right)$

$\vec{F} \cdot \vec{v} = 3t^3 \frac{-2}{t^2} + 6t6 + \frac{2}{t}9t^2 = -6t + 36t + 18t = 48t$

$W = \int_{(2,6,3)}^{(1,12,24)} \vec{F} \cdot d\vec{s} = \int_1^2 \vec{F} \cdot \vec{v} dt = \int_1^2 48t dt = \left[24t^2\right]_1^2 = 24(4 - 1)$

$W = \underline{\quad 72 \quad}$

11. (16 points) Are the following lines parallel, intersecting or skew? If they intersect, find the point of intersection.

a. Line 1: $\vec{r}_1(t) = (t + 2, t - 2, 2t + 1)$

Line 2: $\vec{r}_2(t) = (t + 1, 2t - 6, 2t - 1)$

Solution: The direction vectors, $\vec{v}_1 = (1, 1, 2)$ and $\vec{v}_2 = (1, 2, 2)$, are not multiples of each other. So the lines are not parallel. Since the parameter values may be different at the intersection point, we rewrite the second line as $\vec{r}_2(s) = (s + 1, 2s - 6, 2s - 1)$. We set the x and y components equal to find where the projections intersect in the xy -plane::

$$t + 2 = s + 1 \qquad t - 2 = 2s - 6$$

The first equation says $s = t + 1$. So the second equation says $t - 2 = 2(t + 1) - 6 = 2t - 4$. Or $t = 2$ and so $s = 3$. So the points are

$$\vec{r}_1(2) = (4, 0, 5) \qquad \vec{r}_2(3) = (4, 0, 5)$$

They intersect at $(4, 0, 5)$.

b. Line 1: $\vec{r}_1(t) = (t + 2, t - 2, 2t - 1)$

Line 2: $\vec{r}_2(t) = (t + 1, 2t - 6, 2t - 1)$

Solution: The direction vectors, $\vec{v}_1 = (1, 1, 2)$ and $\vec{v}_2 = (1, 2, 2)$, are not multiples of each other. So the lines are not parallel. Since the parameter values may be different at the intersection point, we rewrite the second line as $\vec{r}_2(s) = (s + 1, 2s - 6, 2s - 1)$. We set the x and y components equal to find where the projections intersect in the xy -plane:

$$t + 2 = s + 1 \qquad t - 2 = 2s - 6$$

The first equation says $s = t + 1$. So the second equation says $t - 2 = 2(t + 1) - 6 = 2t - 4$. Or $t = 2$ and so $s = 3$. So the points are

$$\vec{r}_1(2) = (4, 0, 3) \qquad \vec{r}_2(3) = (4, 0, 5)$$

They do not intersect. They are skew.