

Name _____

MATH 251 Final Exam Version B Fall 2017

Sections 515 Solutions P. Yasskin

Multiple Choice: (5 points each. No part credit.)

1-10	/50	13	/15
11	/5	14	/15
12	/20	Total	/105

1. A wire has the shape of the helix curve $\vec{r}(\theta) = (4 \cos \theta, 4 \sin \theta, 3\theta)$ for $0 \leq \theta \leq \pi$ and has linear density $\delta = 2y$. Find the total mass of the wire.

- a. 80 Correct Choice
- b. 60
- c. 40
- d. 20
- e. 10

Solution: $\delta = 2y = 8 \sin \theta$

$$\vec{v} = (-4 \sin \theta, 4 \cos \theta, 3) \quad |\vec{v}| = \sqrt{16 \sin^2 \theta + 16 \cos^2 \theta + 9} = 5 \quad ds = |\vec{v}| d\theta = 5 d\theta$$

$$M = \int \delta ds = \int_0^\pi 2y |\vec{v}| d\theta = \int_0^\pi 8 \sin \theta 5 d\theta = 40 - \cos \theta \Big|_0^\pi = 80$$

2. A wire has the shape of the helix curve $\vec{r}(\theta) = (4 \cos \theta, 4 \sin \theta, 3\theta)$ for $0 \leq \theta \leq \pi$ and has linear density $\delta = 2y$. Find the y -component of the center of mass of the wire.

- a. 80π
- b. $\frac{1}{80\pi}$
- c. 40π
- d. $\frac{1}{40\pi}$
- e. π Correct Choice

Solution: From the previous problem, $M = 80$.

$$M_y = \int y \delta ds = \int_0^\pi y 2y |\vec{v}| d\theta = \int_0^\pi 32 \sin^2 \theta 5 d\theta = 80 \int_0^\pi (1 + \cos 2\theta) d\theta = 80 \left[\theta + \frac{\sin 2\theta}{2} \right]_0^\pi = 80\pi$$

So the y -component of the center of mass is $\bar{y} = \frac{M_y}{M} = \frac{80\pi}{80} = \pi$

3. The spiral ramp shown at the right may be parametrized by


$$\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, \theta)$$

for $0 \leq r \leq 2$ and $0 \leq \theta \leq 2\pi$.

Find the total mass, if the surface density is

$$\delta = \sqrt{x^2 + y^2}$$

- a. $\frac{2\pi}{3}(5^{3/2} - 1)$ Correct Choice
 b. $\frac{2\pi}{3}5^{3/2}$
 c. $\frac{2\pi}{3}(10^{3/2} - 1)$
 d. $\frac{2\pi}{3}10^{3/2}$
 e. $\frac{2\pi}{3}(10^{3/2} - 5^{3/2})$



Solution: The tangent vectors, normal vector and its length are:

$$\begin{aligned} \vec{e}_r &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \cos \theta & \sin \theta & 0 \\ -r \sin \theta & r \cos \theta & 1 \end{vmatrix} & \vec{N} = \vec{e}_u \times \vec{e}_v = \hat{i}(\sin \theta) - \hat{j}(\cos \theta) + \hat{k}(r \cos^2 \theta + r \sin^2 \theta) \\ \vec{e}_\theta &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -\sin \theta & \cos \theta & 0 \\ r \cos \theta & -r \sin \theta & 1 \end{vmatrix} & = (\sin \theta, -\cos \theta, r) \\ |\vec{N}| &= \sqrt{\sin^2 \theta + \cos^2 \theta + r^2} = \sqrt{1 + r^2} \end{aligned}$$

The density is $\delta = \sqrt{x^2 + y^2} = r$. So the mass is

$$M = \iint \delta dS = \iint \delta |\vec{N}| dr d\theta = \int_0^{2\pi} \int_0^2 r \sqrt{1 + r^2} dr d\theta = 2\pi \left[\frac{(1 + r^2)^{3/2}}{3} \right]_0^2 = \frac{2\pi}{3}(5^{3/2} - 1)$$

4. Consider the spiral ramp described in the previous problem.

Find the flux of the vector field $\vec{F} = (0, 0, z)$ **upward** through the spiral ramp.

- a. $-9\pi^2$
 b. $-4\pi^2$
 c. 0
 d. $4\pi^2$ Correct Choice
 e. $9\pi^2$

Solution: From the previous problem $\vec{N} = (\sin \theta, -\cos \theta, r)$ which is oriented upward.

On the ramp, $\vec{F} = (0, 0, \theta)$. So the flux is

$$\iint \vec{F} \cdot d\vec{S} = \iint \vec{F} \cdot \vec{N} dr d\theta = \int_0^{2\pi} \int_0^2 \theta r dr d\theta = \left[\frac{\theta^2}{2} \right]_0^{2\pi} \left[\frac{r^2}{2} \right]_0^2 = 4\pi^2$$

5. Compute $\int_A^B \vec{F} \cdot d\vec{s}$ for $\vec{F} = (2x + y, x + 2y)$ along the line segment from $A = (2, 1)$ to $B = (1, 3)$.

Hint: Find a scalar potential.

- a. 15
- b. 6 Correct Choice
- c. 0
- d. -6
- e. -15

Solution: $\vec{F} = (2x + y, x + 2y) = \vec{\nabla}f$ for $f = x^2 + xy + y^2$ since $\partial_x f = 2x + y$ and $\partial_y f = x + 2y$.
By the F.T.C.C. $\int \vec{F} \cdot d\vec{s} = \int_{(2,1)}^{(1,3)} \vec{\nabla}f \cdot d\vec{s} = f(1,3) - f(2,1) = (1 + 3 + 9) - (4 + 2 + 1) = 6$

6. Compute $\int_A^B \vec{F} \cdot d\vec{s}$ for $\vec{F} = (-y, x, 3)$ along the helix $\vec{r}(\theta) = (3 \cos \theta, 3 \sin \theta, 4\theta)$ from $A = (3, 0, 0)$ to $B = (3, 0, 8\pi)$.

- a. 0
- b. 40π
- c. 42π Correct Choice
- d. 44π
- e. 46π

Solution: Since $\vec{\nabla} \times \vec{F} = (0, 0, 2) \neq \vec{0}$, there is no scalar potential and we cannot use the F.T.C.C.
So we compute the line integral from its definition:


$$\vec{v} = (-3 \sin \theta, 3 \cos \theta, 4) \quad \vec{F}|_{\vec{r}(\theta)} = (-3 \sin \theta, 3 \cos \theta, 3) \quad \vec{F} \cdot \vec{v} = 9 \sin^2 \theta + 9 \cos^2 \theta + 12 = 21$$

$$\int_A^B \vec{F} \cdot d\vec{s} = \int_0^{2\pi} \vec{F} \cdot \vec{v} d\theta = \int_0^{2\pi} 21 d\theta = 42\pi$$

7. Compute $\oint_{\partial T} (\sin x + 5y) dx + (2x + \cos y) dy$

clockwise around the complete boundary of the triangle shown at the right.

Hint: Use a Theorem.




- a. 12 Correct Choice
- b. 8
- c. 0
- d. -8
- e. -12

Solution: $P = \sin x + 5y$ $Q = 2x + \cos y$ $\partial_x Q - \partial_y P = 2 - 5 = -3$
By Green's Theorem, (There is a minus sign, since we want clockwise.)

$$\oint_{\partial T} \vec{F} \cdot d\vec{s} = - \iint_T (\partial_x Q - \partial_y P) dx dy = - \iint_T -3 dx dy = 3(\text{area}) = 3(4) = 12$$

8. Compute $\oint \vec{F} \cdot d\vec{s}$ for $\vec{F} = (x^2y, 2x^3)$ along the piece of the parabola $y = x^2$ from $(-3, 9)$ to $(3, 9)$ followed by the line segment from $(3, 9)$ back to $(-3, 9)$.



Hint: Use Green's Theorem.

- a. 405
- b. 324 Correct Choice
- c. 162
- d. 81
- e. 0

Solution: $\oint \vec{F} \cdot d\vec{s} = \oint P dx + Q dy$ with $P = x^2y$ and $Q = 2x^3$. By Green's Theorem,
 $\oint \vec{F} \cdot d\vec{s} = \int_{-3}^3 \int_{x^2}^9 (\partial_x Q - \partial_y P) dy dx = \int_{-3}^3 \int_{x^2}^9 (6x^2 - x^2) dy dx = \int_{-3}^3 5x^2 [y]_{x^2}^9 dx$
 $= \int_{-3}^3 (45x^2 - 5x^4) dx = [15x^3 - x^5]_{-3}^3 = 2(15 \cdot 3^3 - 3^5) = 324$

9. Compute $\iint_{\partial C} \vec{F} \cdot d\vec{S}$ over the complete surface of the cylinder $x^2 + y^2 \leq 9$ for $0 \leq z \leq 4$ oriented out from the cylinder for $\vec{F} = (xz, yz, z^2)$.

Hint: Use Gauss' Theorem.

- a. 24π
- b. 36π
- c. 72π
- d. 144π
- e. 288π Correct Choice

Solution: By Gauss' Theorem, $\iint_{\partial C} \vec{F} \cdot d\vec{S} = \iiint_C \vec{\nabla} \cdot \vec{F} dV$ and $\vec{\nabla} \cdot \vec{F} = z + z + 2z = 4z$
 $\iiint_C \vec{\nabla} \cdot \vec{F} dV = \int_0^4 \int_0^{2\pi} \int_0^3 4z r dr d\theta dz = 4 \left[\frac{z^2}{2} \right]_0^4 (2\pi) \left[\frac{r^2}{2} \right]_0^3 = 4(8)(2\pi) \left(\frac{9}{2} \right) = 288\pi$

10. Sketch the region of integration for the integral $\int_0^2 \int_{x^2}^4 x \cos(y^2) dy dx$ in problem (11).

Select its value here:


- a. $\frac{1}{4} \sin 2$
- b. $\frac{1}{4} \sin 4$
- c. $\frac{1}{2} \sin 4$
- d. $\frac{1}{4} \sin 16$ Correct Choice
- e. $\frac{1}{2} \sin 16$

Solution: Reverse the order of integration:

$$\begin{aligned} \int_0^2 \int_{x^2}^4 x \cos(y^2) dy dx &= \int_0^4 \int_0^{\sqrt{y}} x \cos(y^2) dx dy \\ &= \int_0^4 \left[\frac{x^2}{2} \cos(y^2) \right]_{x=0}^{\sqrt{y}} dy = \int_0^4 \frac{y}{2} \cos(y^2) dy \end{aligned}$$

Substitute: $u = y^2 \quad du = 2y dy \quad y dy = \frac{1}{2} du$

$$\begin{aligned} \int_0^2 \int_{x^2}^4 x \cos(y^2) dy dx &= \frac{1}{4} \int \cos(u) du = \frac{1}{4} \sin(u) \\ &= \frac{1}{4} \sin(y^2) \Big|_0^4 = \frac{1}{4} \sin 16 \end{aligned}$$



Work Out: (Points indicated. Part credit possible. Show all work.)


11. (5 points) Sketch the region of integration

for the integral $\int_0^2 \int_{x^2}^4 x \cos(y^2)$.

Shade in the region.

Compute its value in problem (10).

Solution:



12. (20 points) Verify Stokes' Theorem $\iint_P \vec{\nabla} \times \vec{F} \cdot d\vec{S} = \oint_{\partial P} \vec{F} \cdot d\vec{s}$

for the vector field $\vec{F} = (-yz, xz, z^2)$ and the **surface** which is the piece of the paraboloid P given by $z = x^2 + y^2$ between $z = 1$ and $z = 9$ oriented up and in.

Notice that the boundary of P is two circles.



Be sure to check orientations. Use the following steps:

- a. The paraboloid may be parametrized by $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, r^2)$ for $1 \leq r \leq 3$.

$$\vec{e}_r = \begin{pmatrix} \hat{i} \cos \theta \\ \hat{j} \sin \theta \\ 2r \end{pmatrix}$$

$$\vec{e}_\theta = \begin{pmatrix} -r \sin \theta \\ r \cos \theta \\ 0 \end{pmatrix}$$

$$\vec{N} = \vec{e}_r \times \vec{e}_\theta = \hat{i}(-2r^2 \cos \theta) - \hat{j}(2r^2 \sin \theta) + \hat{k}(r \cos^2 \theta + r \sin^2 \theta) = (-2r^2 \cos \theta, -2r^2 \sin \theta, r)$$

\vec{N} has the correct orientation.

$$\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -yz & xz & z^2 \end{vmatrix} = \hat{i}(-x) - \hat{j}(y) + \hat{k}(z+z) = (-x, -y, 2z)$$

$$\vec{\nabla} \times \vec{F} \Big|_{\vec{R}(r, \theta)} = (-r \cos \theta, -r \sin \theta, 2r^2)$$

$$\vec{\nabla} \times \vec{F} \cdot \vec{N} = 2r^3 \cos^2 \theta + 2r^3 \sin^2 \theta + 2r^3 = 4r^3$$

$$\iint_P \vec{\nabla} \times \vec{F} \cdot d\vec{S} = \iint_P \vec{\nabla} \times \vec{F} \cdot \vec{N} dr d\theta = \int_0^{2\pi} \int_1^3 4r^3 dr d\theta = 2\pi [r^4]_{r=1}^3 = 2\pi(81 - 1) = 160\pi$$

Recall $\vec{F} = (-yz, xz, z^2)$

- b. Parametrize the upper circle U and compute the line integral.

$$\vec{r}(\theta) = (3 \cos \theta, 3 \sin \theta, 9)$$

$$\vec{v}(\theta) = (-3 \sin \theta, 3 \cos \theta, 0) \quad \text{oriented correctly counterclockwise}$$

$$\vec{F}|_{\vec{r}(\theta)} = (-27 \sin \theta, 27 \cos \theta, 81)$$

$$\oint_U \vec{F} \cdot d\vec{s} = \int_0^{2\pi} \vec{F} \cdot \vec{v} d\theta = \int_0^{2\pi} 81 \sin^2 \theta + 81 \cos^2 \theta d\theta = \int_0^{2\pi} 81 d\theta = 162\pi$$

- c. Parametrize the lower circle L and compute the line integral.

$$\vec{r}(\theta) = (\cos \theta, \sin \theta, 1)$$

$$\vec{v}(\theta) = (-\sin \theta, \cos \theta, 0) \quad \text{oriented counterclockwise, need clockwise}$$

$$\text{Rev } \vec{v}(\theta) = (\sin \theta, -\cos \theta, 0)$$

$$\vec{F}|_{\vec{r}(\theta)} = (-\sin \theta, \cos \theta, 1)$$

$$\oint_L \vec{F} \cdot d\vec{s} = \int_0^{2\pi} \vec{F} \cdot \vec{v} d\theta = \int_0^{2\pi} -\sin^2 \theta - \cos^2 \theta d\theta = -\int_0^{2\pi} 1 d\theta = -2\pi$$

- d. Combine $\oint_U \vec{F} \cdot d\vec{s}$ and $\oint_L \vec{F} \cdot d\vec{s}$ to get $\oint_{\partial C} \vec{F} \cdot d\vec{s}$

$$\oint_{\partial P} \vec{F} \cdot d\vec{s} = \oint_U \vec{F} \cdot d\vec{s} + \oint_L \vec{F} \cdot d\vec{s} = 162\pi - 2\pi = 160\pi$$

which agrees with part (a).

13. (15 points) (Also replaces Exam 3 #12.)

Find the mass of the solid between the hemispheres

$$z = \sqrt{4 - x^2 - y^2} \quad \text{and} \quad z = \sqrt{9 - x^2 - y^2}$$

for $z \geq 0$ if the density is $\delta = \frac{1}{x^2 + y^2 + z^2}$.



Solution: In spherical coordinates, $2 \leq \rho \leq 3$ and $\delta = \frac{1}{\rho^2}$ and $dV = \rho^2 \sin \phi d\rho d\phi d\theta$.

$$\begin{aligned} M &= \iiint \delta dV = \int_0^{2\pi} \int_0^{\pi/2} \int_2^3 \frac{1}{\rho^2} \rho^2 \sin \phi d\rho d\phi d\theta = 2\pi \left[\rho \right]_2^3 \left[-\cos \phi \right]_0^{\pi/2} \\ &= 2\pi(3 - 2)(-0 - -1) = 2\pi \end{aligned}$$

14. (15 points) (Also replaces Exam 3 #13.)

Find the **centroid** of the **solid** inside

the paraboloid $z = x^2 + y^2$ for $1 \leq z \leq 2$.

Hint: Put the differentials in the order $dr dz d\theta$.



Solution: In cylindrical coordinates, $dV = r dr d\theta dz$ and the paraboloid is $z = r^2$ or $r = \sqrt{z}$.

The volume is:

$$\begin{aligned} V &= \iiint 1 dV = \int_0^{2\pi} \int_1^2 \int_0^{\sqrt{z}} r dr dz d\theta = 2\pi \int_1^2 \left[\frac{r^2}{2} \right]_0^{\sqrt{z}} dz = \pi \int_1^2 z dz = \pi \left[\frac{z^2}{2} \right]_1^2 \\ &= \frac{\pi}{2}(4 - 1) = \frac{3\pi}{2} \end{aligned}$$

The z -moment is:

$$\begin{aligned} V_z &= \iiint z dV = \int_0^{2\pi} \int_1^2 \int_0^{\sqrt{z}} z r dr dz d\theta = 2\pi \int_1^2 z \left[\frac{r^2}{2} \right]_0^{\sqrt{z}} dz \\ &= \pi \int_1^2 z^2 dz = \pi \left[\frac{z^3}{3} \right]_1^2 = \frac{\pi}{3}(8 - 1) = \frac{7\pi}{3} \end{aligned}$$

So the z -component of the centroid is:

$$\bar{z} = \frac{V_z}{V} = \frac{7\pi}{3} \frac{2}{3\pi} = \frac{14}{9} \approx 1.444$$

We know $\bar{x} = \bar{y} = 0$ by symmetry. So the centroid is $(\bar{x}, \bar{y}, \bar{z}) = \left(0, 0, \frac{14}{9}\right)$.