

Name \_\_\_\_\_

MATH 251      Final Exam Version H      Fall 2017

Sections 200      P. Yasskin

Multiple Choice: (5 points each. No part credit.)

|      |     |       |      |
|------|-----|-------|------|
| 1-10 | /50 | 13    | /15  |
| 11   | /5  | 14    | /15  |
| 12   | /20 | Total | /105 |

1. A wire has the shape of the helix curve  $\vec{r}(\theta) = (4 \cos \theta, 4 \sin \theta, 3\theta)$  for  $0 \leq \theta \leq \pi$  and has linear density  $\delta = 2y$ . Find the total mass of the wire.

- a. 80
- b. 60
- c. 40
- d. 20
- e. 10

2. A wire has the shape of the helix curve  $\vec{r}(\theta) = (4 \cos \theta, 4 \sin \theta, 3\theta)$  for  $0 \leq \theta \leq \pi$  and has linear density  $\delta = 2y$ . Find the  $y$ -component of the center of mass of the wire.

- a.  $80\pi$
- b.  $\frac{1}{80\pi}$
- c.  $40\pi$
- d.  $\frac{1}{40\pi}$
- e.  $\pi$

3. The spiral ramp shown at the right may be parametrized by

$$\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, \theta)$$

for  $0 \leq r \leq 2$  and  $0 \leq \theta \leq 2\pi$ .

Find the total mass, if the surface density is

$$\delta = \sqrt{x^2 + y^2}$$

- a.  $\frac{2\pi}{3}(5^{3/2} - 1)$
- b.  $\frac{2\pi}{3}5^{3/2}$
- c.  $\frac{2\pi}{3}(10^{3/2} - 1)$
- d.  $\frac{2\pi}{3}10^{3/2}$
- e.  $\frac{2\pi}{3}(10^{3/2} - 5^{3/2})$



4. Consider the spiral ramp described in the previous problem.

Find the flux of the vector field  $\vec{F} = (0, 0, z)$  **upward** through the spiral ramp.

- a.  $-9\pi^2$
- b.  $-4\pi^2$
- c. 0
- d.  $4\pi^2$
- e.  $9\pi^2$

5. Compute  $\int_A^B \vec{F} \cdot d\vec{s}$  for  $\vec{F} = (2x + y, x + 2y)$  along the line segment from  $A = (2, 1)$  to  $B = (1, 3)$ .

**Hint:** Find a scalar potential.

- a. 15
- b. 6
- c. 0
- d. -6
- e. -15


6. Compute  $\int_A^B \vec{F} \cdot d\vec{s}$  for  $\vec{F} = (-y, x, 3)$  along the helix  $\vec{r}(\theta) = (3 \cos \theta, 3 \sin \theta, 4\theta)$  from  $A = (3, 0, 0)$  to  $B = (3, 0, 8\pi)$ .

- a. 0
- b.  $40\pi$
- c.  $42\pi$
- d.  $44\pi$
- e.  $46\pi$

7. Compute  $\oint_{\partial T} (\sin x + 5y) dx + (2x + \cos y) dy$


**clockwise** around the complete boundary of the triangle shown at the right.

**Hint:** Use a Theorem.



- a. 12
- b. 8
- c. 0
- d. -8
- e. -12

8. Compute  $\oint \vec{F} \cdot d\vec{n}$  for  $\vec{F} = (2x^3, -x^2y)$  along the piece of the parabola  $y = x^2$  from  $(-3, 9)$  to  $(3, 9)$  followed by the line segment from  $(3, 9)$  back to  $(-3, 9)$ .



**Hint:** Use the 2D Gauss' Theorem.

- a. 405  
 b. 324  
 c. 162  
 d. 81  
 e. 0
9. Compute  $\iint_{\partial C} \vec{F} \cdot d\vec{S}$  over the complete surface of the cylinder  $x^2 + y^2 \leq 9$  for  $0 \leq z \leq 4$  oriented out from the cylinder for  $\vec{F} = (xz, yz, z^2)$ .

**Hint:** Use Gauss' Theorem.

- a.  $24\pi$   
 b.  $36\pi$   
 c.  $72\pi$   
 d.  $144\pi$   
 e.  $288\pi$

10. Sketch the region of integration for the integral  $\int_0^2 \int_{x^2}^4 x \cos(y^2) dy dx$  in problem (11).

Select its value here:

- a.  $\frac{1}{4} \sin 2$
- b.  $\frac{1}{4} \sin 4$
- c.  $\frac{1}{2} \sin 4$
- d.  $\frac{1}{4} \sin 16$
- e.  $\frac{1}{2} \sin 16$

---


Work Out: (Points indicated. Part credit possible. Show all work.)

11. (5 points) Sketch the region of integration

for the integral  $\int_0^2 \int_{x^2}^4 x \cos(y^2)$ .

Shade in the region.

Compute its value in problem (10).



12. (20 points) Verify Stokes' Theorem  $\iint_P \vec{\nabla} \times \vec{F} \cdot d\vec{S} = \oint_{\partial P} \vec{F} \cdot d\vec{s}$

for the vector field  $\vec{F} = (-yz, xz, z^2)$  and the **surface** which is the piece of the paraboloid  $P$  given by  $z = x^2 + y^2$  between  $z = 1$  and  $z = 9$  oriented up and in.

Notice that the boundary of  $P$  is two circles.

Be sure to check orientations. Use the following steps:

- a. The paraboloid may be parametrized by  $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, r^2)$  for  $1 \leq r \leq 3$ .



$$\vec{e}_r =$$

$$\vec{e}_\theta =$$

$$\vec{N} =$$

(Check the orientation)

$$\vec{\nabla} \times \vec{F} =$$

$$\vec{\nabla} \times \vec{F} \Big|_{\vec{R}(r, \theta)} =$$

$$\vec{\nabla} \times \vec{F} \cdot \vec{N} =$$

$$\iint_P \vec{\nabla} \times \vec{F} \cdot d\vec{S} =$$

Recall  $\vec{F} = (-yz, xz, z^2)$

- b. Parametrize the upper circle  $U$  and compute the line integral.

$$\vec{r}(\theta) =$$

$$\vec{v}(\theta) =$$

(Check the orientation)

$$\vec{F}|_{\vec{r}(\theta)} =$$

$$\oint_U \vec{F} \cdot d\vec{s} =$$

- c. Parametrize the lower circle  $L$  and compute the line integral.

$$\vec{r}(\theta) =$$

$$\vec{v}(\theta) =$$

(Check the orientation)

$$\vec{F}|_{\vec{r}(\theta)} =$$

$$\oint_L \vec{F} \cdot d\vec{s} =$$

- d. Combine  $\oint_U \vec{F} \cdot d\vec{s}$  and  $\oint_L \vec{F} \cdot d\vec{s}$  to get  $\oint_{\partial C} \vec{F} \cdot d\vec{s}$ .

13. (15 points) (Also replaces Exam 3 #12.)

Find the mass of the solid between the hemispheres

$$z = \sqrt{4 - x^2 - y^2} \quad \text{and} \quad z = \sqrt{9 - x^2 - y^2}$$

for  $z \geq 0$  if the density is  $\delta = \frac{1}{x^2 + y^2 + z^2}$ .



14. (15 points) (Also replaces Exam 3 #13.)

Find the **centroid** of the **solid** inside

the paraboloid  $z = x^2 + y^2$  for  $1 \leq z \leq 2$ .

**Hint:** Put the differentials in the order  $dr dz d\theta$ .

