

MATH 304

Linear Algebra

**Lecture 36:**

**Complex eigenvalues and eigenvectors.  
Symmetric and orthogonal matrices.**

## Complex numbers

$\mathbb{C}$ : complex numbers.

Complex number:  $z = x + iy,$

where  $x, y \in \mathbb{R}$  and  $i^2 = -1$ .

$i = \sqrt{-1}$ : imaginary unit

Alternative notation:  $z = x + yi$ .

$x$  = real part of  $z$ ,

$iy$  = imaginary part of  $z$

$y = 0 \implies z = x$  (real number)

$x = 0 \implies z = iy$  (purely imaginary number)

We add, subtract, and multiply complex numbers as polynomials in  $i$  (but keep in mind that  $i^2 = -1$ ).

If  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$ , then

$$z_1 + z_2 = (x_1 + x_2) + i(y_1 + y_2),$$

$$z_1 - z_2 = (x_1 - x_2) + i(y_1 - y_2),$$

$$z_1 z_2 = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1).$$

Given  $z = x + iy$ , the **complex conjugate** of  $z$  is  $\bar{z} = x - iy$ . The **modulus** of  $z$  is  $|z| = \sqrt{x^2 + y^2}$ .

$$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2 = |z|^2.$$

$$z^{-1} = \frac{\bar{z}}{|z|^2}, \quad (x + iy)^{-1} = \frac{x - iy}{x^2 + y^2}.$$

## Complex exponentials

*Definition.* For any  $z \in \mathbb{C}$  let

$$e^z = 1 + z + \frac{z^2}{2!} + \cdots + \frac{z^n}{n!} + \cdots$$

*Remark.* A sequence of complex numbers  $z_1 = x_1 + iy_1$ ,  $z_2 = x_2 + iy_2$ ,  $\dots$  converges to  $z = x + iy$  if  $x_n \rightarrow x$  and  $y_n \rightarrow y$  as  $n \rightarrow \infty$ .

**Theorem 1** If  $z = x + iy$ ,  $x, y \in \mathbb{R}$ , then

$$e^z = e^x(\cos y + i \sin y).$$

In particular,  $e^{i\phi} = \cos \phi + i \sin \phi$ ,  $\phi \in \mathbb{R}$ .

**Theorem 2**  $e^{z+w} = e^z \cdot e^w$  for all  $z, w \in \mathbb{C}$ .

**Proposition**  $e^{i\phi} = \cos \phi + i \sin \phi$  for all  $\phi \in \mathbb{R}$ .

*Proof:* 
$$e^{i\phi} = 1 + i\phi + \frac{(i\phi)^2}{2!} + \dots + \frac{(i\phi)^n}{n!} + \dots$$

The sequence  $1, i, i^2, i^3, \dots, i^n, \dots$  is periodic:



$$\underbrace{1, i, -1, -i, 1, i, -1, -i, \dots}$$

It follows that

$$\begin{aligned} e^{i\phi} &= 1 - \frac{\phi^2}{2!} + \frac{\phi^4}{4!} - \dots + (-1)^k \frac{\phi^{2k}}{(2k)!} + \dots \\ &+ i \left( \phi - \frac{\phi^3}{3!} + \frac{\phi^5}{5!} - \dots + (-1)^k \frac{\phi^{2k+1}}{(2k+1)!} + \dots \right) \\ &= \cos \phi + i \sin \phi. \end{aligned}$$

## Geometric representation

Any complex number  $z = x + iy$  is represented by the vector/point  $(x, y) \in \mathbb{R}^2$ .



$$x = r \cos \phi, \quad y = r \sin \phi \implies z = r(\cos \phi + i \sin \phi) = re^{i\phi}$$

If  $z_1 = r_1 e^{i\phi_1}$  and  $z_2 = r_2 e^{i\phi_2}$ , then

$$z_1 z_2 = r_1 r_2 e^{i(\phi_1 + \phi_2)}, \quad z_1 / z_2 = (r_1 / r_2) e^{i(\phi_1 - \phi_2)}.$$

## Fundamental Theorem of Algebra

Any polynomial of degree  $n \geq 1$ , with complex coefficients, has exactly  $n$  roots (counting with multiplicities).

Equivalently, if

$$p(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0,$$

where  $a_i \in \mathbb{C}$  and  $a_n \neq 0$ , then there exist complex numbers  $z_1, z_2, \dots, z_n$  such that

$$p(z) = a_n (z - z_1)(z - z_2) \cdots (z - z_n).$$

## Complex eigenvalues and eigenvectors

*Example.*  $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ .  $\det(A - \lambda I) = \lambda^2 + 1$ .

Characteristic roots:  $\lambda_1 = i$  and  $\lambda_2 = -i$ .

Associated eigenvectors:  $\mathbf{v}_1 = \begin{pmatrix} 1 \\ -i \end{pmatrix}$  and  $\mathbf{v}_2 = \begin{pmatrix} 1 \\ i \end{pmatrix}$ .

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ -i \end{pmatrix} = \begin{pmatrix} i \\ 1 \end{pmatrix} = i \begin{pmatrix} 1 \\ -i \end{pmatrix},$$

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ i \end{pmatrix} = \begin{pmatrix} -i \\ 1 \end{pmatrix} = -i \begin{pmatrix} 1 \\ i \end{pmatrix}.$$

$\mathbf{v}_1, \mathbf{v}_2$  is a basis of eigenvectors. *In which space?*



## Complexification

Instead of the real vector space  $\mathbb{R}^2$ , we consider a *complex vector space*  $\mathbb{C}^2$  (all complex numbers are admissible as scalars).

The linear operator  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,  $f(\mathbf{x}) = A\mathbf{x}$  is extended to a *complex linear operator*  $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ ,  $F(\mathbf{x}) = A\mathbf{x}$ .

The vectors  $\mathbf{v}_1 = (1, -i)$  and  $\mathbf{v}_2 = (1, i)$  form a basis for  $\mathbb{C}^2$ .

$\mathbb{C}^2$  is also a real vector space (of real dimension 4). The standard real basis for  $\mathbb{C}^2$  is  $\mathbf{e}_1 = (1, 0)$ ,  $\mathbf{e}_2 = (0, 1)$ ,  $i\mathbf{e}_1 = (i, 0)$ ,  $i\mathbf{e}_2 = (0, i)$ . The matrix of the operator  $F$  with respect to this basis has block structure  $\begin{pmatrix} A & O \\ O & A \end{pmatrix}$ .

## Dot product of complex vectors

Dot product of real vectors

$$\mathbf{x} = (x_1, \dots, x_n), \mathbf{y} = (y_1, \dots, y_n) \in \mathbb{R}^n:$$

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \dots + x_ny_n.$$

Dot product of complex vectors

$$\mathbf{x} = (x_1, \dots, x_n), \mathbf{y} = (y_1, \dots, y_n) \in \mathbb{C}^n:$$

$$\mathbf{x} \cdot \mathbf{y} = x_1\bar{y}_1 + x_2\bar{y}_2 + \dots + x_n\bar{y}_n.$$

If  $z = r + it$  ( $t, s \in \mathbb{R}$ ) then  $\bar{z} = r - it$ ,  
 $z\bar{z} = r^2 + t^2 = |z|^2$ .

Hence  $\mathbf{x} \cdot \mathbf{x} = |x_1|^2 + |x_2|^2 + \dots + |x_n|^2 \geq 0$ .

Also,  $\mathbf{x} \cdot \mathbf{x} = 0$  if and only if  $\mathbf{x} = \mathbf{0}$ .

The norm is defined by  $\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}}$ .

## Normal matrices

*Definition.* An  $n \times n$  matrix  $A$  is called

- **symmetric** if  $A^T = A$ ;
- **orthogonal** if  $AA^T = A^T A = I$ , i.e.,  $A^T = A^{-1}$ ;
- **normal** if  $AA^T = A^T A$ .

**Theorem** Let  $A$  be an  $n \times n$  matrix with real entries. Then

- (a)  $A$  is normal  $\iff$  there exists an orthonormal basis for  $\mathbb{C}^n$  consisting of eigenvectors of  $A$ ;
- (b)  $A$  is symmetric  $\iff$  there exists an orthonormal basis for  $\mathbb{R}^n$  consisting of eigenvectors of  $A$ .

*Example.*  $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 1 \end{pmatrix}.$

- $A$  is symmetric.
- $A$  has three eigenvalues: 0, 2, and 3.
- Associated eigenvectors are  $\mathbf{v}_1 = (-1, 0, 1)$ ,  $\mathbf{v}_2 = (1, 0, 1)$ , and  $\mathbf{v}_3 = (0, 1, 0)$ , respectively.
- Vectors  $\frac{1}{\sqrt{2}}\mathbf{v}_1$ ,  $\frac{1}{\sqrt{2}}\mathbf{v}_2$ ,  $\mathbf{v}_3$  form an orthonormal basis for  $\mathbb{R}^3$ .

**Theorem** Suppose  $A$  is a normal matrix. Then for any  $\mathbf{x} \in \mathbb{C}^n$  and  $\lambda \in \mathbb{C}$  one has

$$A\mathbf{x} = \lambda\mathbf{x} \iff A^T\mathbf{x} = \bar{\lambda}\mathbf{x}.$$

Thus any normal matrix  $A$  shares with  $A^T$  all real eigenvalues and the corresponding eigenvectors.

Also,  $A\mathbf{x} = \lambda\mathbf{x} \iff A\bar{\mathbf{x}} = \bar{\lambda}\bar{\mathbf{x}}$  for any matrix  $A$  with real entries.

**Corollary** All eigenvalues  $\lambda$  of a symmetric matrix are real ( $\bar{\lambda} = \lambda$ ). All eigenvalues  $\lambda$  of an orthogonal matrix satisfy  $\bar{\lambda} = \lambda^{-1} \iff |\lambda| = 1$ .

## Why are orthogonal matrices called so?

**Theorem** Given an  $n \times n$  matrix  $A$ , the following conditions are equivalent:

- (i)  $A$  is orthogonal:  $A^T = A^{-1}$ ;
- (ii) columns of  $A$  form an orthonormal basis for  $\mathbb{R}^n$ ;
- (iii) rows of  $A$  form an orthonormal basis for  $\mathbb{R}^n$ .

*Proof:* Entries of the matrix  $A^T A$  are dot products of columns of  $A$ . Entries of  $AA^T$  are dot products of rows of  $A$ .

Thus an orthogonal matrix is the transition matrix from one orthonormal basis to another.

*Example.*  $A_\phi = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}.$

- $A_\phi A_\psi = A_{\phi+\psi}$
- $A_\phi^{-1} = A_{-\phi} = A_\phi^T$
- $A_\phi$  is orthogonal
- $\det(A_\phi - \lambda I) = (\cos \phi - \lambda)^2 + \sin^2 \phi.$
- Eigenvalues:  $\lambda_1 = \cos \phi + i \sin \phi = e^{i\phi},$   
 $\lambda_2 = \cos \phi - i \sin \phi = e^{-i\phi}.$
- Associated eigenvectors:  $\mathbf{v}_1 = (1, -i),$   
 $\mathbf{v}_2 = (1, i).$
- Vectors  $\frac{1}{\sqrt{2}}\mathbf{v}_1$  and  $\frac{1}{\sqrt{2}}\mathbf{v}_2$  form an orthonormal basis for  $\mathbb{C}^2.$

Consider a linear operator  $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $L(\mathbf{x}) = A\mathbf{x}$ , where  $A$  is an  $n \times n$  orthogonal matrix.

**Theorem** There exists an orthonormal basis for  $\mathbb{R}^n$  such that the matrix of  $L$  relative to this basis has a diagonal block structure

$$\begin{pmatrix} D_{\pm 1} & O & \dots & O \\ O & R_1 & \dots & O \\ \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & R_k \end{pmatrix},$$

where  $D_{\pm 1}$  is a diagonal matrix whose diagonal entries are equal to 1 or  $-1$ , and

$$R_j = \begin{pmatrix} \cos \phi_j & -\sin \phi_j \\ \sin \phi_j & \cos \phi_j \end{pmatrix}, \quad \phi_j \in \mathbb{R}.$$