

MATH 415  
Modern Algebra I

**Lecture 9:**  
**Direct product of groups.**  
**Factor groups.**

## Direct product of groups

Given nonempty sets  $G$  and  $H$ , the Cartesian product  $G \times H$  is the set of all ordered pairs  $(g, h)$  such that  $g \in G$  and  $h \in H$ . Suppose  $*$  is a binary operation on  $G$  and  $\star$  is a binary operation on  $H$ . Then we can define a binary operation  $\bullet$  on  $G \times H$  by

$$(g_1, h_1) \bullet (g_2, h_2) = (g_1 * g_2, h_1 \star h_2).$$

**Proposition 1** The operation  $\bullet$  is fully (resp. uniquely, well) defined if and only if both  $*$  and  $\star$  are.

**Proposition 2** The operation  $\bullet$  is associative if and only if both  $*$  and  $\star$  are associative.

**Proposition 3** A pair  $(e_G, e_H)$  is the identity element in  $G \times H$  if and only if  $e_G$  is the identity element in  $G$  and  $e_H$  is the identity element in  $H$ .

**Proposition 4**  $(g', h') = (g, h)^{-1}$  in  $G \times H$  if and only if  $g' = g^{-1}$  in  $G$  and  $h' = h^{-1}$  in  $H$ .

## Direct product of groups

Given nonempty sets  $G$  and  $H$ , the Cartesian product  $G \times H$  is the set of all ordered pairs  $(g, h)$  such that  $g \in G$  and  $h \in H$ . Suppose  $*$  is a binary operation on  $G$  and  $\star$  is a binary operation on  $H$ . Then we can define a binary operation  $\bullet$  on  $G \times H$  by

$$(g_1, h_1) \bullet (g_2, h_2) = (g_1 * g_2, h_1 \star h_2).$$

**Theorem** The set  $G \times H$  with the operation  $\bullet$  is a group if and only if both  $(G, *)$  and  $(H, \star)$  are groups.

The group  $G \times H$  is called the **direct product** of the groups  $G$  and  $H$ . Usually the same notation (multiplicative or additive) is used for all three groups:

$$\begin{aligned}(g_1, h_1)(g_2, h_2) &= (g_1 g_2, h_1 h_2) \text{ or} \\ (g_1, h_1) + (g_2, h_2) &= (g_1 + g_2, h_1 + h_2).\end{aligned}$$

Similarly, we can define the direct product  $G_1 \times G_2 \times \cdots \times G_n$  of any finite collection of groups  $G_1, G_2, \dots, G_n$ .

*Example.*  $\mathbb{Z}_2 \times \mathbb{Z}_3$  (with  $+_2$  in  $\mathbb{Z}_2$  and  $+_3$  in  $\mathbb{Z}_3$ ).

The group consists of 6 elements. It is abelian since  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  are both abelian. The identity element is  $(0, 0)$ .

Let  $g = (1, 1)$ . Then  $2g = g + g = (0, 2)$ ,  $3g = (1, 0)$ ,  $4g = (0, 1)$ ,  $5g = (1, 2)$ , and  $6g = (0, 0)$ . It follows that  $\mathbb{Z}_2 \times \mathbb{Z}_3$  is a cyclic group,  $\mathbb{Z}_2 \times \mathbb{Z}_3 = \langle g \rangle$ .

**Theorem** If  $g$  has finite order in a group  $G$  and  $h$  has finite order in a group  $H$ , then  $(g, h)$  has finite order in  $G \times H$  equal to  $\text{lcm}(o(g), o(h))$ .

**Theorem** The direct product of nontrivial cyclic groups is cyclic if and only if they are all finite and their orders are pairwise coprime.

For example, groups  $\mathbb{Z}_3 \times \mathbb{Z}_5$ ,  $\mathbb{Z}_4 \times \mathbb{Z}_{15}$ , and  $\mathbb{Z}_2 \times \mathbb{Z}_5 \times \mathbb{Z}_7$  are cyclic while groups  $\mathbb{Z}_4 \times \mathbb{Z}_6$ ,  $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_3 \times \mathbb{Z}$ , and  $\mathbb{Z} \times \mathbb{Z}$  are not.

## Factor space

Let  $X$  be a nonempty set and  $\sim$  be an equivalence relation on  $X$ . Given an element  $x \in X$ , the **equivalence class** of  $x$ , denoted  $[x]_{\sim}$  or simply  $[x]$ , is the set of all elements of  $X$  that are **equivalent** (i.e., related by  $\sim$ ) to  $x$ :

$$[x]_{\sim} = \{y \in X \mid y \sim x\}.$$

**Theorem** Equivalence classes of the relation  $\sim$  form a partition of the set  $X$ .

The set of all equivalence classes of  $\sim$  is denoted  $X/\sim$  and called the **factor space** (or **quotient space**) of  $X$  by the relation  $\sim$ .

In the case when the set  $X$  carries some structure (algebraic, geometric, analytic, etc.), this structure may (or may not) induce an analogous structure on the factor space  $X/\sim$ .

## Examples of factor spaces

- $X = G$ , a group;  $x \sim y$  if and only if  $x \in yH$ , where  $H$  is a fixed subgroup.

Equivalence class of an element  $g \in G$  is a left coset of the subgroup  $H$ ,  $[g]_{\sim} = gH$ . The factor space  $G/\sim$  is the set of all left cosets of  $H$  in  $G$ . It is usually denoted  $G/H$ .

- $X = G$ , a group;  $x \sim y$  if and only if  $x \in Hy$ , where  $H$  is a fixed subgroup.

Equivalence class of an element  $g \in G$  is a right coset of the subgroup  $H$ ,  $[g]_{\sim} = Hg$ . The factor space  $G/\sim$  is the set of all right cosets of  $H$  in  $G$ . It is often denoted  $H \backslash G$ .

- $X = G$ , a group;  $x \sim y$  if and only if  $x \in KyH = \{kyh : h \in H, k \in K\}$ , where  $H$  and  $K$  are fixed subgroups.

In this example,  $[g]_{\sim} = KgH$  (a **double coset**). The factor space  $G/\sim$  is usually denoted  $K \backslash G/H$ .

## Factor group

Let  $G$  be a nonempty set with a binary operation  $*$ . Given an equivalence relation  $\sim$  on  $G$ , we say that the relation  $\sim$  is **compatible** with the operation  $*$  if for any  $g_1, g_2, h_1, h_2 \in G$ ,

$$g_1 \sim g_2 \text{ and } h_1 \sim h_2 \implies g_1 * h_1 \sim g_2 * h_2.$$

If this is the case, we can define an operation on the factor space  $G/\sim$  by  $[g] \star [h] = [g * h]$  for all  $g, h \in G$ .

Compatibility is required so that the operation  $\star$  is defined uniquely: if  $[g'] = [g]$  and  $[h'] = [h]$  then  $[g' * h'] = [g * h]$ .

If the operation  $*$  is associative (resp. commutative), then so is  $\star$ . If  $e$  is the identity element for  $*$ , then its equivalence class  $[e]$  is the identity element for  $\star$ . If  $h = g^{-1}$  in  $(G, *)$ , then  $[h] = [g]^{-1}$  in  $(G/\sim, \star)$ .

Thus, if  $(G, *)$  is a group then  $(G/\sim, \star)$  is also a group called the **factor group** (or **quotient group**). Moreover, if the group  $(G, *)$  is abelian then so is  $(G/\sim, \star)$ .

**Question.** When is an equivalence relation  $\sim$  on a group  $G$  compatible with the operation?

Let  $G$  be a group and assume that an equivalence relation  $\sim$  on  $G$  is compatible with the operation (so that the factor space  $G/\sim$  is also the factor group). For simplicity, let us use multiplicative notation.

**Lemma 1** The equivalence class of the identity element is a subgroup of  $G$ .

*Proof.* Let  $H = [e]_{\sim}$  be the equivalence class of the identity element  $e$ . We need to show that **(i)**  $e \in H$ , **(ii)**  $h_1, h_2 \in H \implies h_1 h_2 \in H$ , and **(iii)**  $h \in H \implies h^{-1} \in H$ .

By reflexivity,  $e \sim e$ . Hence  $e \in H$ . Further, if  $h_1, h_2 \in H$ , then  $h_1 \sim e$  and  $h_2 \sim e$ . By compatibility,  $h_1 h_2 \sim ee = e$  so that  $h_1 h_2 \in H$ . Next, if  $h \in H$  then  $h \sim e$ . Also,  $h^{-1} \sim h^{-1}$ . By compatibility,  $hh^{-1} \sim eh^{-1}$ , that is,  $e \sim h^{-1}$ . By symmetry,  $h^{-1} \sim e$  so that  $h^{-1} \in H$ .



**Lemma 2** Each equivalence class is a left coset of the subgroup  $H = [e]_{\sim}$ .

*Proof.* We need to prove that  $[g]_{\sim} = gH$  for all  $g \in G$ . We are going to show that  $gH \subset [g]_{\sim}$  and  $[g]_{\sim} \subset gH$ .

Suppose  $a \in gH$ , that is,  $a = gh$  for some  $h \in H$ . Then  $g \sim g$  and  $h \sim e$ , which implies that  $gh \sim ge = g$ . Hence  $a \in [g]_{\sim}$ . Conversely, suppose  $a \in [g]_{\sim}$ . We have  $a = ea = (gg^{-1})a = g(g^{-1}a)$ . Since  $g^{-1} \sim g^{-1}$  and  $a \sim g$ , it follows that  $g^{-1}a \sim g^{-1}g = e$ . Hence  $g^{-1}a \in H$  so that  $a = g(g^{-1}a) \in gH$ .

**Lemma 3** Each equivalence class is a right coset of the subgroup  $H = [e]_{\sim}$ .

*Proof.* Analogous to the proof of Lemma 2.

**Definition.** A subgroup  $H$  of a group  $G$  is called **normal** if  $gH = Hg$  for all  $g \in G$ , that is, each left coset of  $H$  is also a right coset. *Notation:*  $H \triangleleft G$  or  $H \trianglelefteq G$ .

## Factor group

**Question.** When is an equivalence relation  $\sim$  on a group  $G$  compatible with the operation?

**Theorem** Assume that the factor space  $G/\sim$  is also a factor group. Then

- (i)  $H = [e]_{\sim}$ , the equivalence class of the identity element, is a subgroup of  $G$ ,
- (ii)  $[g]_{\sim} = gH$  for all  $g \in G$ ,
- (iii)  $G/\sim = G/H$ ,
- (iv) the subgroup  $H$  is **normal**, which means that  $gH = Hg$  for all  $g \in G$ .

**Theorem** If  $H$  is a normal subgroup of a group  $G$ , then  $G/H$  is a factor group.

## Alternative construction of the factor group

Suppose  $G$  is a group (with multiplicative notation). For any  $X, Y \subset G$  let  $XY = \{xy \mid x \in X, y \in Y\}$ . This “multiplication of sets” is a well-defined operation on  $\mathcal{P}(G)$ , the set of all subsets of  $G$ . The operation is associative:  $(XY)Z = X(YZ)$  for any sets  $X, Y, Z \subset G$ . Indeed,

$$(XY)Z = \{(xy)z \mid x \in X, y \in Y, z \in Z\},$$
$$X(YZ) = \{x(yz) \mid x \in X, y \in Y, z \in Z\}.$$

**Proposition** If  $H$  is a normal subgroup of  $G$ , then for all  $a, b \in G$  we have  $(aH)(bH) = (ab)H$  in the sense of the above definition.

## Alternative construction of the factor group

Suppose  $G$  is a group (with multiplicative notation). For any sets  $X, Y \subset G$  let  $XY = \{xy \mid x \in X, y \in Y\}$ .

**Proposition** If  $H$  is a normal subgroup of  $G$ , then for all  $a, b \in G$  we have  $(aH)(bH) = (ab)H$  in the sense of the above definition.

*Proof.* In terms of multiplication of sets, any coset  $gH$  can be written as  $\{g\}H$ . Therefore  $(aH)(bH) = (\{a\}H)(\{b\}H)$ . By associativity, this is the same as  $\{a\}(H\{b\})H$ . Now  $H\{b\}$  is the right coset  $Hb$ . Since the subgroup  $H$  is normal, we have  $Hb = bH = \{b\}H$ . Again by associativity,

$$(aH)(bH) = \{a\}(\{b\}H)H = (\{a\}\{b\})(HH).$$

Clearly,  $\{a\}\{b\} = \{ab\}$ . It remains to show that  $HH = H$ . Indeed,  $HH \subset H$  since the subgroup  $H$  is closed under the operation. Conversely,  $H = \{e\}H \subset HH$ .